

Methodological Characteristics and Future Directions for Plyometric Jump Training Research: A Scoping Review

Ramírez-Campillo, Rodrigo; Álvarez, Cristian; García-Hermoso, Antonio; Ramírez-Vélez, Robinson; Gentil, Paulo; Asadi, Abbas; Chaabene, Helmi; Moran, Jason; Meylan, César; Garcia de Alcaraz, Antonio; Sánchez, Javier; Nakamura, Fábio Yuzo; Granacher, Urs; Kraemer, William; Izquierdo, Mikel

Published in:
Sports Medicine

Publication date:
2018

The re-use license for this item is:
CC BY-NC-ND

This document version is the:
Peer reviewed version

The final published version is available direct from the publisher website at:
[10.1007/s40279-018-0870-z](https://doi.org/10.1007/s40279-018-0870-z)

[Find this output at Hartpury Pure](#)

Citation for published version (APA):

Ramírez-Campillo, R., Álvarez, C., García-Hermoso, A., Ramírez-Vélez, R., Gentil, P., Asadi, A., Chaabene, H., Moran, J., Meylan, C., Garcia de Alcaraz, A., Sánchez, J., Nakamura, F. Y., Granacher, U., Kraemer, W., & Izquierdo, M. (2018). Methodological Characteristics and Future Directions for Plyometric Jump Training Research: A Scoping Review. *Sports Medicine*, 48(5), 1059-1081. <https://doi.org/10.1007/s40279-018-0870-z>

Methodological Characteristics and Future Directions for Plyometric Jump Training Research: A Scoping Review.

Short Title: Scoping Review of Plyometric Jump Training Studies.

This is a post-peer-review, pre-copyedit version of an article published in Sports Medicine. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s40279-018-0870-z>

Rodrigo Ramirez-Campillo¹ (r.ramirez@ulagos.cl; ORCID: 0000-0003-2035-3279), Cristian Álvarez¹ (cristian.alvarez@ulagos.cl), Antonio García-Hermoso² (antonio.garcia.h@usach.cl; ORCID: 0000-0002-1397-7182), Robinson Ramírez-Vélez³ (robin640@hotmail.com; ORCID: 0000-0003-3075-6960), Paulo Gentil⁴ (paulogentil@hotmail.com; ORCID: 0000-0003-2459-4977), Abbas Asadi⁵ (abbas_asadi1175@yahoo.com), Helmi Chaabene^{6,7} (chaabanehelmi@hotmail.fr; ORCID: 0000-0001-7812-7931), Jason Moran⁸ (jason.moran@hartpury.ac.uk), Cesar Meylan^{9,10} (cmeylan@csipacific.ca), Antonio García de Alcaraz¹¹ (antoniogadealse@gmail.com; ORCID: 0000-0001-8210-4118), Javier Sanchez-Sanchez¹² (jsanchezsa@upsa.es; ORCID: 0000-0003-4585-247X), Fabio Y. Nakamura^{13,14} (fabioy_nakamura@yahoo.com.br), Urs Granacher⁶ (urs.granacher@uni-potsdam.de; ORCID: 0000-0002-7095-813X), William Kraemer¹⁵ (kraemer.44@osu.edu), Mikel Izquierdo¹⁶ (mikel.izquierdo@gmail.com; ORCID: 0000-0002-1506-4272).

¹. Department of Physical Activity Sciences. Research Nucleus in Health, Physical Activity and Sport. Universidad de Los Lagos. Osorno. Chile.

². Laboratorio de Ciencias de la Actividad Física, el Deporte y la Salud, Facultad de Ciencias Médicas, Universidad de Santiago de Chile, USACH, Santiago, Chile.

³. Centro de Estudios para la Medición de la Actividad Física “CEMA”, Escuela de Medicina y Ciencias de la Salud, Universidad del Rosario, Bogotá D.C, Colombia.

⁴. Faculdade de Educação Física e Dança, Universidade Federal de Goiás, Goiânia, Brasil.

⁵. Department of Physical Education and Sport Sciences, Payame Noor University, Tehran, Iran.

⁶. Division of Training and Movement Sciences, Research Focus Cognition Sciences, University of Potsdam, Am Neuen Palais 10, 14469 Potsdam, Germany

⁷. High Institute of Sports and Physical Education, Kef, University of Jendouba, Tunisia

⁸. Department of Sport, University Centre Hartpury, University of the West of England.

- ⁹. Canadian Sport Institute Pacific, Vancouver, Canada.
- ¹⁰. Canadian Soccer Association, Ottawa, Canada.
- ¹¹. LFE Research Group, Faculty of Physical Activity and Sports Sciences, Universidad Politécnica de Madrid, Madrid, Spain.
- ¹². Research Group Planning and Assessment of Training and Athletic Performance. Pontifical University of Salamanca, Salamanca, Spain.
- ¹³. The College of Healthcare Sciences, James Cook University, Queensland, Australia.
- ¹⁴. Department of Medicine and Aging Sciences, “G. d’Annunzio” University of Chieti-Pescara, Italy.
- ¹⁵. Neuroscience/Neuromuscular Human Research Laboratory, Department of Human Sciences, The Ohio State University
- ¹⁶. Department of Health Sciences, Public University of Navarre, Navarra, Spain.

Corresponding Author

Department of Health Sciences

Public University of Navarre (Navarra) SPAIN

Campus of Tudela

Av. de Tarazona s/n. 31500 Tudela (Navarra) SPAIN

Tel + 34 948 417876

mikel.izquierdo@gmail.com

No sources of funding were used to assist in the preparation of the present review.

The authors declare that they have no conflict of interest relevant to the content of this review.

ABSTRACT

Recently, there has been a proliferation of published articles on the effect of plyometric jump training including several review articles and meta-analyses. However, these types of research articles are generally of narrow scope. Furthermore, methodological limitations among studies (e.g., lack of active/passive control groups) prevent the generalization of results and these factors need to be addressed by researchers. On that basis, the aims of this scoping review were to i) characterize the main elements of plyometric jump training studies (e.g., training protocols), and ii) provide future directions for research. From 648 potentially relevant articles, 242 were eligible for inclusion in this review. The main issues identified related to: an insufficient number of studies conducted in females, youths and individual sports (~24.0%, ~37.0% and ~12.0% of overall studies, respectively); insufficient reporting of effect size values and training prescription (~34.0% and ~55.0% of overall studies, respectively); studies missing an active/passive control group and randomization (~40.0% and ~20.0% of overall studies, respectively). Furthermore, plyometric jump training was often combined with other training methods and added to participants' daily training routines (~47.0% and ~39.0% of overall studies, respectively), thus distorting conclusions on its independent effects. Additionally, most studies lasted no longer than 7 weeks. In future, researchers are advised to conduct plyometric training studies of high methodological quality (e.g., randomized controlled trials). More research is needed in females, youth, and individual sports. Finally, the identification of specific dose-response relationships following plyometric training are needed to specifically tailor intervention programs, particularly in the long-term.

Key Points

- Recently, there has been a proliferation of published articles on the effect of plyometric jump training in various populations.
- There has been relatively few studies conducted in females, youths and individual sports, with insufficient reporting of effect size values and training prescription, studies missing an active/passive control group and randomization, and most studies lasting no longer than 7 weeks.
- In the future, researchers are advised to conduct plyometric training studies of high methodological quality (e.g., randomized controlled trials) with more research needed on specific dose-response relationships, particularly in the long-term.

1. INTRODUCTION

Efficient use of the stretch-shortening cycle is a key factor in jump performance with the accumulation of elastic energy in a muscle action facilitating greater mechanical work in subsequent actions [1-3]. Jump exercises are an effective means of physical conditioning for the promotion of health, injury prevention and skill-related measures of athletic performance [4-6]. Jump training is commonly associated with plyometric training and, in particular, with drills that stress the musculotendinous unit [1, 2]. In this sense, plyometric jump training involves the utilization of different types of jumping movements [7-9] that are incorporated into comprehensive strength training programs [5, 10, 11]. Numerous plyometric jump training articles have been published in peer-reviewed journals in recent years [12]. However, between the years of 2000 and 2017, scientific publications on plyometric jump training have increased 25 fold compared with any previous period [12]. This rapid expansion has been driven by the technology-related advancement of appropriate measuring devices [13, 14] and an increased awareness amongst scientists of plyometrics' potential benefits for athletic performance [15], health (e.g., improve the bone mineral density) [16-19], injury prevention [20], rehabilitation [21] and application to special populations [22-24].

Several high-quality reviews related to plyometric jump training have been published [25-28, 15, 29, 30], with a particular focus on the effects on athletic performance (e.g., agility, muscular strength, sprint, and jump performance), as well as in specific populations (e.g., male youth athletes) and various sports (e.g., soccer). Such reviews and meta-analyses generally adopt strict inclusion criteria for studies' eligibility [31, 25, 27, 32], thus reducing the limitation of potential bias. However, the application of such rigorous inclusion criteria tends to limit the number of studies in these reviews, resulting in the capturing of only some of the large number of articles that have been published. Accordingly, conclusions from reviews and meta-analyses may thus be limited [33, 34] in their facility to assess the methodological gaps and limitations among the articles that have been published.

Given the above observations, the aims of this scoping review article were to: i) characterize the main methodological features of the body of literature relating to plyometric jump training studies (e.g., participants' characteristics; training protocols), and ii) recommend future directions for plyometric jump training research. Information gathered for this scoping review could be useful for sports scientists, practitioners, and, ultimately, athletes, as it may help in the understanding of the methodological gaps and limitations in the current plyometric jump training literature. Additionally, this review may inspire the design of more high-quality studies in the field, resulting in fewer sources of bias and greater generalizability to relevant populations.

2. METHODS

2.1. Search Strategy

A literature search for published studies on the PubMed database (<https://www.ncbi.nlm.nih.gov/pubmed/?term=plyometric+training>) [35] was performed up to April 17th 2017. This search was conducted using the term ‘plyometric training’.

Considering the broad international scope of the current review, we searched for articles in all languages (e.g., fifty-four) available on the PubMed database. Only original research was included. However, to reduce the chances of excluding potentially relevant studies, seventy-two article types available on the PubMed database were included during the initial search, with no restriction for journal categories, including those found on MEDLINE. In addition, no age, sex, or publication date criteria were imposed during the initial search. Default values of the PubMed database search engine were used in search fields, PubMed commons, text availability (e.g., abstract, free full text, and full text), and participants. Manual data checking was performed to increase the precision of data collection from relevant studies.

2.2. Retrieved Articles Selection

After an initial search, an account was created in the National Center for Biotechnology Information (<https://www.ncbi.nlm.nih.gov>). Through this account, the lead investigator received automatically generated emails for updates regarding the search terms used. These updates were received on a daily basis and studies were eligible for inclusion until the initiation of manuscript preparation on July 1st, 2017.

To facilitate the inclusion of a sufficiently high number of articles, many different study designs were considered. This is in line with the principles of scoping reviews [36-38], including those in the field of resistance training [5, 39]. This resulted in the accumulation of a rather broad body of research and, consequently, the reviewing of studies with different levels of quality and research questions. Several limitations were systematically identified according to standard data extraction criteria (e.g., PEDro scale) and applied to selected articles for quality assessment [40]. This is detailed in subsequent sections of this article. In line with the aforementioned principles of scoping reviews [36-38, 5, 39], both non-randomized and non-controlled studies were included [41].

Inclusion criteria. Included studies were those that incorporated pre- and post-intervention performance testing with ≥ 6 plyometric jump training sessions over a period ≥ 2 weeks, including plyometric jump drills as a primary component (individually; or embedded in a wider training program). Articles that used plyometric jump training with added resistance were also included. Conference contributions were considered if full-text articles were available.

Exclusion criteria. Articles were excluded if they were cross-sectional (transversal), a review, or a training-related study without plyometric jump training. Also excluded were retrospective studies, prospective studies, studies in which the use of jump exercises was not clearly described, studies for which only the abstract was available, case reports, studies with ambiguous study protocols, non-human investigations, special communications, repeated bout effect interventions, repeated references, letters to the editor, invited commentaries, errata, overtraining studies and detraining studies. In the case of detraining studies, if there was a training period prior to a detraining period, the study was considered for inclusion.

2.3. Data Extraction

From potentially relevant retrieved articles, generic information (e.g., author's name; journal name; year) and abstracts were saved for analyses. Two investigators (RRC and CA) independently processed all data with one extracting and the other verifying. Based on previous recommendations for improving searching in PubMed [42], suggestions for plyometric jump training searches [32, 27, 29, 25] and expert opinion on methodological gaps and limitations of plyometric jump training studies, several data items were considered for extraction. A detailed outline of each of these items is provided in table 1. The items were grouped into three broad subjects for results presentation and discussion purposes: i) main general characteristics; ii) participants' characteristics; iii) key elements of plyometric jump training design.

TABLE 1 NEAR HERE

2.4. Eligible Articles

From 648 articles, 406 (62.6%) were excluded (Figure 1) leaving 242 (37.4%) eligible for data extraction [43-106, 22, 107-144, 24, 145-220, 20, 221-280] (Table 2).

FIGURE 1 NEAR HERE

TABLE 2 NEAR HERE

3. RESULTS

The 242 eligible articles appeared in 47 different journals. The majority (n=241) were published in English, with a clear exponential increase in number in recent years (Figure 2).

FIGURE 2 NEAR HERE

3.1. General Characteristics

From eligible articles, 54.5% were classified as being insufficiently described (see table 1 for further information regarding the identification of treatment description quality).

Note: the remaining results are derived from table 2 (unless stated otherwise).

From the 242 eligible studies, 40.5% did not incorporate an active/passive control group and 19.4% did not consider any form of randomization. However, the number of eligible articles that have incorporated a randomized and controlled design have increased from 1987 to 2017, comprising up to 52.0% of relevant literature in recent years, compared to 33.3% previously. The number of participants included per study averaged 15.9 participants. From the total number of reported dependent variables (n=3982), 14.6% were deemed not clearly or quantitatively available (e.g., presented in graphical form only), whilst changes reported as effect sizes were available for only 34.0% of these (data not shown in this scoping review).

3.2. Participants' Characteristics

Most studies included male participants (57.2%) whilst 23.9% were female participants and 16.6% were a mixture of both sexes. Of note, the sex of participants was not clearly reported in 2.3% of the studies. Most of the studies included participants with a mean age ≥ 18 years old and 36.7% included youth participants, with groups' mean age equal to 19.6 years old. Participants' mean body mass, stature, and body mass index were 65.6 kg, 170.4 cm, and 22.6 kg·m⁻², respectively.

The physical performance/playing level of participants was high (e.g., professional/elite athletes) in 14.9% of the studies, low (e.g., participants under medical treatment) in 8.3% of the studies, and not-clearly reported in 5.3% of the studies. Participants with moderate (e.g., non-elite/non-professional athletes) or normal (e.g., recreational athletes) physical performance or playing level featured in 46.3% of studies. Participants with mixed physical performance or playing level were included in 25.2% of studies (see table 1 for a detailed depiction of physical performance/playing level classification). With regard to the type of sport practiced, 44.2% included athletes from team sports (mostly soccer players [31.1%]), 12.0% undertook individual sports (mostly endurance runners [6.7%]), 8.7% played both individual and team sports whilst 30.1% were non-competitive (e.g., recreationally) athletes. In 5.0% of studies, this information was not clearly reported. Twelve percent of studies included participants with experience of plyometric jump training, 46.7% without experience, and 41.3% did not clearly report these details. Amongst the

included studies, 14.0% focused on training during the pre-season period of the season, 28.5% during the in-season, and 5.8% during the off-season. In 17.8% of studies, this information was not clearly reported whilst 33.9% of the studies did not include participants who were competitive athletes or who were engaged in a year-round training plan.

3.3. Key Elements of Plyometric Jump Training Design

The types of plyometric jump drills were not clearly reported in 2.9% of the studies whilst 72.3% used a mixture ≥ 2 jump types. In addition, 24.8% employed one type of jump only. Drop jumps were included in 32.1% of studies with a combination of heights used in 17.4% and individualized prescription of heights used in 1.3% of studies. Box heights for drop jumps ranged from 10 to 110 cm.

Underfoot surface type was not clearly reported in 64.2% of studies. A proportion of 7% of studies reported either clay, wood, athletic track, cement or gymnasium-type floors. With regard to softer surfaces, 2.2% of studies reported the use of athletic mats, 7.8% utilised grass surfaces and 4.0% incorporated aquatic, elastic or sand-based surfaces. A mixture of surface types were used in 7.0% of the studies (e.g., inclined-flat; land-aquatic). In 7.0% of studies, plyometric jump training was completed using special equipment which included force plates, unstable surfaces, sledge apparatus, or similar commercially available apparatus, other than boxes or barriers. With regard to training loads, 5.2% of studies did not clearly report the number of jumps completed during the intervention period. Amongst those that did report the number of jumps, a wide range of values was observed, with 250 jumps per week as a mean. However, values varied according to training design (e.g., duration).

Plyometric jump training was added to the habitual training of participants in 38.5% of studies. In 20.5% of studies, part of the habitual training strategy of participants was replaced with plyometric jump training; however, 22.1% of the studies did not clearly report the approach taken on this issue. Approximately 18.9% of the studies recruited subjects who were not involved in habitual physical training. Plyometric jump training was combined with other methods of training as part of an intervention in 47.1% of studies but no clear information was identified in 2.5% of the studies. Most of the studies that applied a combination of plyometric jump training with other methods used resistance (29.7%), speed (14.3%), and agility (8.0%) training. Other training methods that were used in combination with plyometric jump training were sport-specific, balance, electrostimulation, stretching, core, footwork, endurance, coordination, and high-intensity interval training methods. Most commonly, two or more of these methods were used in combination with plyometric jump training.

Training duration ranged from 2 to 96 weeks. Most studies applied ≤ 7 weeks of training (mode), with a mean number of 8.3 weeks observed. However, there was high variability for this statistic with a standard deviation of 7.2 weeks. With regard to training frequency, 5.6%, 50.6%, 28.5%, 1.6%, and 0.4% of the studies used one, two, three, four, and six sessions per week, respectively. From the included studies, 12.9% used a combination of training frequencies, more commonly two and three sessions per week. Some groups were submitted to a range of training frequencies ranging from zero up to five sessions per week. Plyometric jump training intensity was not clearly reported for 42.0% of the included studies while 52.0% reported it as maximal using criteria such as height, distance, reactive strength index, optimal power, percentage of one repetition maximum, time, voluntary effort, velocity, rate of execution, force, or a mixture of these. Training intensity was submaximal in 6.0% of studies and this was quantified using similar measures such as percentage of one repetition maximum, height, distance, velocity, and rating of perceived exertion.

A proportion of 17.7% of the studies did not feature progressive overload as an element of programme design and 6.7% did not present clear information relating to this factor. Individual overload techniques (e.g., volume-based; technique-based; intensity-based) were applied in 34.2% of the studies whereas 41.4% of studies used a mixed model of progression, combining two or three individual overload techniques during the training program. Information relating to the tapering of training was not reported in 11.1% of the studies. Conversely, 15.5% applied a tapering strategy but 73.4% did not present clear information with respect to this factor.

The rest time between sets and/or exercises was not clearly reported for 42.4% of the studies. The rest interval extended from zero to 600-s, with a mean of 119-s, a standard deviation of 80-s, and a mode of 120-s. With regard to the rest period between plyometric jump repetitions, 79.3% of the studies did not clearly specify the interval (if any) between plyometric jumps. For those that reported the duration, this ranged from 4-s to 120-s, a mean of 13-s, a standard deviation of 8-s, and a mode of 15-s. The rest period between training sessions was not clearly reported in 45.6% of the studies. Among those studies that reported this value, 48 and 72 hours were the most common rest period durations reported, with intervals ranging from 24 hours to 168 hours.

4. DISCUSSION

The aims of this scoping review article were to: i) characterize the main methodological features of the literature relating to plyometric jump training studies; ii) provide recommendations for future plyometric jump training research. The main results allowed a comprehensive characterization of the main methodological features of the literature of plyometric jump training

studies. In the following paragraphs a discussion is provided on the implications of the identified elements and future directions for plyometric jump training research (table 3).

TABLE 3 NEAR HERE

4.1. Main General Characteristics

From the 242 eligible articles ~51% were not described in sufficient detail meaning that their findings could not be effectively leveraged for future research or practice. Accordingly, just under 50% of the studies in this review demonstrate the quality of description required to leverage their findings for future research and practice. Examples of insufficiently described study features relate to the omission of one or more basic treatment descriptors such as training duration, frequency, intensity, exercise type and number of prescribed sets and repetitions. Moreover, around 40% of the included studies did not incorporate a control group *per se* and approximately 25% were non-randomized or did not clearly report information relating to this design element. However, randomization and controlled design implementation has become more common, increasing from 33.3% to 52% of published articles in recent years. This seems to indicate that the growing number of published articles is mirrored by a concurrent improvement in the methodological design of plyometric jump training interventions. With regard to the reporting of results, around 15.0% of dependent variable changes were not clearly described, often being presented in graphical form only whilst effect size changes were reported for just 34% of dependent variables. Given the growing consensus with regard to the importance of effect sizes in intervention studies [281], investigators should consider not only the null-hypothesis test but also the magnitude-based inference analysis [282]. In relation to this issue, p-values could be argued to be pernicious given the binary nature of their associated conclusions which relate to the statistical terms ‘*significant*’ and ‘*non-significant*’ [283]. In addition, the adequate classification of participants categories according to training experience and level (e.g., professional athlete) should be considered by researchers [284].

Although some limitations are difficult to address (e.g., subjects’ blinding), investigators should strive to conduct more randomized and controlled studies whilst also being mindful that the nature of the control condition will depend heavily on the applied setting in which the study is conducted. For example, researchers must consider the use of physically active control groups in an athletic setting and whilst passive controls are more appropriate in a clinical setting. When using a control group is not possible, a washout period or a cross over design might serve as a potential alternative in a field setting though this may be a suboptimal approach in youth athletes who are physically maturing at a fast rate. Studies should include adequate sample sizes which can sustain high rates of participant attrition and descriptions of utilised methods should be detailed to the degree that

studies could be replicated without need for recourse. If necessary, study details should be included in a supplemental file. Results should be comprehensively reported and should include effect sizes and associated inferences.

4.2. Participants' Characteristics

As in similar research fields [285], relatively few (<25%) plyometric jump training studies included females, either adult or youths. Most of the included studies recruited participants with a mean age of around 18 years and less than 37% of the included studies involved youth groups. Amongst the studies that included youth participants, biological age was reported only in around 37% of study groups, an important research gap previously identified [5]. This limitation is compounded by the utilization of different measures of maturity across studies making it difficult to compare results. This could be viewed as a critical limitation among plyometric jump training interventions performed in youths because biological age seems to influence adaptations to plyometric jump training interventions [25]. Similarly, only one group with a mean age of over 65 years was identified. This is concerning given that modified jump exercises for older adults may be a good alternative [22] to *conventional* jumps. Regarding anthropometric data, most studies were conducted in healthy-weight participants. Some studies included overweight and obese participants [62, 61] whilst others reported changes in participants' body composition [94, 132, 159, 175, 210] or anatomical adaptations related to reduced injury risk (e.g., tendon hypertrophy) [286-288, 139]. Although usually seen as a training method to induce neural-related adaptations, plyometric jump training deserves further consideration as a training strategy to induce musculoskeletal [4] and body composition adaptations.

Only around 15.0% of the included studies were conducted in participants with a high-level of athletic performance or sport playing level. This may be due to the unwillingness of professional or elite athletes' coaches to modify their training schedule. Similarly, although plyometric jump training may have potential relevance as a rehabilitation therapy [107, 180, 226, 116, 146, 178, 61, 62, 96, 24, 201], only around 8% of the included studies were conducted in participants with a low-level of athletic performance or playing level. Moreover, the moderating effect of sports practice is relatively uncertain amongst individual sports as only around 12% of the included studies have been conducted in athletes who were engaged in such activities. Nevertheless, it does seem that plyometric jump training offers beneficial adaptations to participants with different athletic performance or playing levels and in various sports disciplines. Regarding plyometric jump training experience, although some reviews and meta-analyses reported no effects of previous experience on sprint [29], maximal strength [30], vertical jump [28], and agility [27] adaptations after a plyometric jump training intervention, others reported the opposite [26]. Moreover, as of yet, no controlled comparative research has been conducted and around 41.0% of included studies did not clearly report on this issue. However, it

is reasonable to expect that previous experience with jump training (e.g., long-jumpers) could impact on the degree of adaptation in high-level athletes because their ceiling for adaptation could be lower.

Cross-sectional studies have shown that plyometric jump training is applied regularly during athletes' competitive season [289-297]. Its application on a year-round basis might reduce the occurrence of injuries, most notably among youth athletes [298]. However, based on this review, the identification of cases in which the application of plyometric jump training was conducted in a year-round fashion proved difficult. Moreover, only 48.0% of the included studies applied plyometric jump training at some point of the participant's season (e.g., in-season), without a long-term follow-up in order to compare its effects during other moments of the season (e.g., pre-season).

4.3. Key Elements of Plyometric Jump Training Design

To assess the specific effects of plyometric jump training in isolation with its effects as a part of a wider training program, researchers must clarify whether interventions are carried out with, without, or in place of, other forms of physical activity [189]. Despite this, nearly 22.0% of the included studies did not report if plyometric training was added *to* an existing training program or if it *replaced* part of a program. Relatedly, 39% of the included studies reported that plyometric training was added to a program whilst ~47.0% of the included studies reported that it was added to another training method as part of an intervention (resistance, sprint, agility, balance, electrostimulation, stretching, core, footwork, endurance, coordination and high-intensity interval training). Considering this, researchers should aim to strengthen their methodological approaches by clarifying if the effects of interventions occur because of plyometric jump training only or are due to a combination with another training method within or without a regular program of physical preparation. In this way, researchers could consider replacing part of athletes' habitual training with plyometric jump training, avoiding the introduction of other training methods to prevent the distortion of results.

Most of the included studies combined two or more types of jump drills, which, from a practical perspective, is considered to be a sound approach to optimizing adaptations [91, 88]. Despite this, from a research perspective, the prescription of multiple jump types within a single intervention could confound the independent effect of a single jump type, for example, a drop jump. Accordingly, more research is needed to assess the specific effects of different plyometric jumps types with researchers being encouraged to apply programs that have little, if any, variation in the prescribed jump drill. Although ~25.0% of the included studies sought to assess the effect of a single type of plyometric jump, these studies did not compare the effect of one type of jump versus another. Moreover, very few investigations have compared the effects of different jump intensities (e.g., drop jump heights) [246] or the effects of intensity distribution (e.g., polarized plyometric training), whilst only three included studies

individualized the prescription of jump training. Although it is beyond the scope of the current review to provide a detailed description of the dependent variables analyzed amongst included articles, it is worth mentioning that only three studies analyzed the effect of plyometric jump training on injury rates [20, 222, 249]. Relatedly, injury incidence was not associated with participation in plyometric jump training programs, including those that incorporated drop jumps, thus underlining the apparent safety of plyometric jump training.

The type of training surface used for plyometric jump training may affect the speed of the stretch-shortening cycle that is performed (e.g., fast vs. slow), thus implying different biomechanical and physiological effects [299, 300] and possibly different adaptations. However, the surface type was not clearly reported for most (64.2%) of the included studies. The use of soft surfaces was reported by less than 15% of the included studies and, in most cases, their use was related to the specific nature of athletes' competition surface (e.g., grass surface for soccer players) [280]. Plyometric jump training was completed using special equipment in around 7% of the included studies. The utilization of special equipment to aid plyometric jump training adaptations is still a matter of debate, especially with regard to the usefulness of unstable surfaces compared to stable surfaces [301]. Noticeably, only one study quantified (e.g., by means of the restitution coefficient) the hardness of training surface [138]. It is particularly noteworthy that authors do not typically acknowledge this factor which could be justifiably considered a key aspect of plyometric program configuration.

Different physiological and physical performance traits (e.g., agility; speed) exhibit different temporal responses to plyometric jump training [27, 30, 28, 29, 22, 302]. In this sense, training duration should be adjusted to the type of trait that is being targeted by the coach or athlete. However, most of the included studies lasted just 7 weeks or less. Therefore, the long-term effects of plyometric jump training are difficult to characterize based on evidence from the current body of available literature. In addition, the applied training volume varied widely among studies with optimal values still yet to be determined. Although plyometric jump training volume recommendations have been provided for some physical performance traits [29, 30, 28], a general lack of consensus might partially explain the wide variation in current opinion. Training volume should be closely considered for both athletic performance enhancement and injury prevention purposes [303] as relevant doses could differ [272]. The weekly distribution of plyometric training frequency of included studies ranged from one to six sessions per week, with around 79% of the included studies applying 2 to 3 sessions per week. Some studies compared the effects of different training frequencies and revealed that this factor might not be as important as the overall training volume [45, 204, 211]. This corroborates the findings of some meta-analyses [27, 304] yet contrasts with those of others [28, 30, 29].

Plyometric jump training intensity has been defined as the amount of stress placed on involved muscles and connective tissue and joints during a given exercise [305]. Commonly, athletes perform drop jumps from progressively increasing heights for a more intense training stimulus [306, 307, 145, 308]. Indeed, it has been shown that there is greater muscle activity in drop jumps from a 60-cm box height than from a 40-cm [309] or 20-cm box height [306]. However, these assertions are not without opposition [310-314]. Considering this, although training intensity is a key factor to acknowledge in exercise prescription [11], plyometric jump training lacks consensus on optimal markers of intensity. Relatedly, the intensity of plyometric jump training was not clearly reported in around 42% of included studies. Moreover, when reported, in several cases the description of how intensity was achieved was difficult to derive or was reported using highly varied criteria such as height, distance, reactive strength index, optimal power, percentage of one repetition maximum, time, voluntary effort, velocity, rate of execution, force, rating of perceived exertion or a mixture of these factors.

Progressive overload implicates a gradual increase of the stress placed upon the body during exercise training and is considered to be one of the fundamental principles of training [11, 87]. However, 23.2% of included studies did not incorporate an element of progressive overload or failed to report clear information relating to this factor. While around 33% of the included studies used a one-dimensional form of overload (e.g., volume-, technique- or intensity-based), approximately 41% used a mixed model of progression. However, the difference between the effects of various progression models adopted in the literature still requires elucidation. After an overload period, the application of a training taper has been identified as a key strategy [315-318]. This generally consists of reducing the training volume towards the end of a program or prior to a competition (or assessment) period [317]. However, only 15.5% of the included studies incorporated a taper strategy. Moreover, around 85% of the included studies did not apply a taper or failed to report it. On that basis, the effects of a taper as part of a plyometric training protocol need to be further examined.

Although evidence has been provided on the acute effects of rest time between plyometric jump repetitions [319] and sets/exercises [320], studies which directly investigate this are scarce [102]. Indeed, the rest time between repetitions and sets/exercises was not clearly reported for 79.3% and 42.4% of the included studies, respectively. When this information was reported, there were wide variations in the rest intervals utilized. Considering that the required rest period may depend on factors such as the duration of the effort [321, 322] and participants' ages [323], the wide variety of rest intervals used among the included studies may be explained by the various type of jump drills and the wide age range of participants. Regarding rest between training sessions, the most commonly used period was 48 to 72 hours. Even though anecdotal reports recommend that plyometric jump training sessions should not be performed on consecutive days [324], there is a lack of experimental studies that have addressed

this issue [120]. Factors such as training experience, habituation to plyometric jump drills, age and intensity could be considered when rest periods between sessions are assessed [4].

Aside from several strengths (e.g., the number of analyzed studies; the broad spectrum of items extracted from studies), this review has some limitations. Factors such as the volume of plyometric jumps completed as a warm-up, the training attendance rate, and the coach-to-trainee ratio during interventions were not considered as review criteria. In addition, given the broad aim of this review, only the term '*plyometric training*' was used in the search strategy.

5. CONCLUSION

From 648 potentially relevant articles, 242 were eligible for data extraction with 26 variables from each allowing a comprehensive characterization of the main methodological features of the literature relating to plyometric jump training. This was based on the initial identification of ~6292 elements through database searching. The implications of study design features for future directions in plyometric jump training research were described. To our knowledge, this one of the most extensive and comprehensive review studies conducted so far regarding plyometric jump training.

For investigators working in the field of plyometric jump training, a primary goal is to mobilize institutional, political and government bodies, in addition to coaches and athletes, to increase awareness about the importance of this type of physical training activity. Benefits derived from plyometric jump training may include improvement of neuromuscular, metabolic, and cardiovascular markers as well as body composition and proxies of athletic performance. This can be achieved in both females and males as well as in pediatric and geriatric populations, sedentary participants and competitive athletes, both in preventive and rehabilitative therapy settings. Investigators involved in the relevant field should strive to increase funds available for research and should also look to foster international and multicenter collaboration toward a better understanding of the benefits that can be derived from this type of training strategy. This can be achieved through the promotion of a higher quality of research output. Overall, through this scoping review, several methodological elements have been recognized that should be addressed in future studies.

6. REFERENCES

1. Taube W, Leukel C, Gollhofer A. How neurons make us jump: the neural control of stretch-shortening cycle movements. *Exerc Sport Sci Rev.* 2012;40(2):106-15. doi:10.1097/JES.0b013e31824138da.

2. Komi PV, Gollhofer A. Stretch reflex can have an important role in force enhancement during SSC-exercise. *Journal of Applied Biomechanics*. 1997;13:451-9.
3. Marey M, Demeny M. Locomotion humaine, mecanisme du saut. *C R Acad Sci (Paris)*. 1885;101:489-94.
4. Markovic G, Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sports Med*. 2010;40(10):859-95. doi:10.2165/11318370-000000000-00000.
5. Granacher U, Lesinski M, Busch D, Muehlbauer T, Prieske O, Puta C et al. Effects of resistance training in youth athletes on muscular fitness and athletic performance: a conceptual model for long-term athlete development. *Frontiers in physiology*. 2016;7:164. doi:10.3389/fphys.2016.00164.
6. Cormie P, McGuigan MR, Newton RU. Developing maximal neuromuscular power: part 2 - training considerations for improving maximal power production. *Sports Med*. 2011;41(2):125-46. doi:10.2165/11538500-000000000-00000.
7. Duda M. Plyometrics: A Legitimate Form of Power Training? *Phys Sportsmed*. 1988;16(3):212-8. doi:10.1080/00913847.1988.11709466.
8. Faigenbaum A, Chu D. *Plyometric Training for Children and Adolescents*. Indianapolis, IN: American College of Sports Medicine. 2017.
9. Sands WA, Wurth JJ, Hewitt JK. Speed and Agility Training. In: (NSCA) TNSaCAs, editor. *Basics of Strength and Conditioning manual*. 2012.
10. Behm DG, Faigenbaum AD, Falk B, Klenstrup P. Canadian Society for Exercise Physiology position paper: resistance training in children and adolescents. *Appl Physiol Nutr Metab*. 2008;33(3):547-61. doi:10.1139/H08-020.
11. American College of Sports Medicine. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc*. 2009;41(3):687-708. doi:10.1249/MSS.0b013e3181915670.
12. PubMed NTNCfBI. Plyometric training. 2017. <https://www.ncbi.nlm.nih.gov/pubmed>. Accessed April 17 2017.
13. Gallardo-Fuentes F, Gallardo-Fuentes J, Ramirez-Campillo R, Balsalobre-Fernandez C, Martinez C, Caniuqueo A et al. Intersession and Intrasession Reliability and Validity of the My Jump App for Measuring Different Jump Actions in Trained Male and Female Athletes. *J Strength Cond Res*. 2016;30(7):2049-56. doi:10.1519/jsc.0000000000001304.
14. Balsalobre-Fernandez C, Glaister M, Lockett RA. The validity and reliability of an iPhone app for measuring vertical jump performance. *J Sports Sci*. 2015;33(15):1574-9. doi:10.1080/02640414.2014.996184.
15. Markovic G. Does plyometric training improve vertical jump height? A meta-analytical review. *Br J Sports Med*. 2007;41(6):349-55; discussion 55. doi:10.1136/bjsm.2007.035113.
16. Gomez-Bruton A, Matute-Llorente A, Gonzalez-Aguero A, Casajus JA, Vicente-Rodriguez G. Plyometric exercise and bone health in children and adolescents: a systematic review. *World J Pediatr*. 2017;13(2):112-21. doi:10.1007/s12519-016-0076-0.

17. Xu J, Lombardi G, Jiao W, Banfi G. Effects of Exercise on Bone Status in Female Subjects, from Young Girls to Postmenopausal Women: An Overview of Systematic Reviews and Meta-Analyses. *Sports Med.* 2016;46(8):1165-82. doi:10.1007/s40279-016-0494-0.
18. Zhao R, Zhao M, Zhang L. Efficiency of jumping exercise in improving bone mineral density among premenopausal women: a meta-analysis. *Sports Med.* 2014;44(10):1393-402. doi:10.1007/s40279-014-0220-8.
19. Zribi A, Zouch M, Chaari H, Bouajina E, Ben Nasr H, Zaouali M et al. Short-Term Lower-Body Plyometric Training Improves Whole-Body BMC, Bone Metabolic Markers, and Physical Fitness in Early Pubertal Male Basketball Players. *Pediatr Exerc Sci.* 2014;26(1):22-32. doi:10.1123/pes.2013-0053.
20. Myrick S. Injury prevention and performance enhancement: a training program for basketball. *Conn Med.* 2007;71(1):5-8.
21. Chmielewski TL, Myer GD, Kauffman D, Tillman SM. Plyometric exercise in the rehabilitation of athletes: physiological responses and clinical application. *J Orthop Sports Phys Ther.* 2006;36(5):308-19. doi:10.2519/jospt.2006.2013.
22. Piirainen JM, Cronin NJ, Avela J, Linnamo V. Effects of plyometric and pneumatic explosive strength training on neuromuscular function and dynamic balance control in 60-70year old males. *J Electromyogr Kinesiol.* 2014;24(2):246-52. doi:10.1016/j.jelekin.2014.01.010.
23. Johnson BA, Salzberg CL, Stevenson DA. Effects of a plyometric training program for 3 children with neurofibromatosis type 1. *Pediatr Phys Ther.* 2012;24(2):199-208. doi:10.1097/PEP.0b013e31824d30ee.
24. Gonzalez-Aguero A, Vicente-Rodriguez G, Gomez-Cabello A, Ara I, Moreno LA, Casajus JA. A 21-week bone deposition promoting exercise programme increases bone mass in young people with Down syndrome. *Dev Med Child Neurol.* 2012;54(6):552-6. doi:10.1111/j.1469-8749.2012.04262.x.
25. Moran JJ, Sandercock GR, Ramirez-Campillo R, Meylan CM, Collison JA, Parry DA. Age-Related Variation in Male Youth Athletes' Countermovement Jump After Plyometric Training: A Meta-Analysis of Controlled Trials. *J Strength Cond Res.* 2017;31(2):552-65. doi:10.1519/JSC.0000000000001444.
26. Stojanovic E, Ristic V, McMaster DT, Milanovic Z. Effect of Plyometric Training on Vertical Jump Performance in Female Athletes: A Systematic Review and Meta-Analysis. *Sports Med.* 2017;47(5):975-86. doi:10.1007/s40279-016-0634-6.
27. Asadi A, Arazi H, Young WB, Saez de Villarreal E. The Effects of Plyometric Training on Change-of-Direction Ability: A Meta-Analysis. *Int J Sports Physiol Perform.* 2016;11(5):563-73. doi:10.1123/ijsp.2015-0694.
28. de Villarreal ES, Kellis E, Kraemer WJ, Izquierdo M. Determining variables of plyometric training for improving vertical jump height performance: a meta-analysis. *J Strength Cond Res.* 2009;23(2):495-506. doi:10.1519/JSC.0b013e318196b7c6.
29. Saez de Villarreal E, Requena B, Cronin JB. The effects of plyometric training on sprint performance: a meta-analysis. *J Strength Cond Res.* 2012;26(2):575-84. doi:10.1519/JSC.0b013e318220fd03.

30. Saez-Saez de Villarreal E, Requena B, Newton RU. Does plyometric training improve strength performance? A meta-analysis. *J Sci Med Sport*. 2010;13(5):513-22. doi:10.1016/j.jsams.2009.08.005.
31. Bedoya AA, Miltenberger MR, Lopez RM. Plyometric training effects on athletic performance in youth soccer athletes: A systematic review. *J Strength Cond Res*. 2015;29(8):2351-60. doi:10.1519/JSC.0000000000000877.
32. Behm DG, Young JD, Whitten JHD, Reid JC, Quigley PJ, Low J et al. Effectiveness of traditional strength vs. power training on muscle strength, power and speed with youth: a systematic review and meta-analysis. *Frontiers in physiology*. 2017;8(423). doi:10.3389/fphys.2017.00423.
33. Greco T, Zangrillo A, Biondi-Zoccai G, Landoni G. Meta-analysis: pitfalls and hints. *Heart, lung and vessels*. 2013;5(4):219-25.
34. Gentil P, Arruda A, Souza D, Giessing J, Paoli A, Fisher J et al. Is there any practical application of meta-analytical results in strength training? *Frontiers in physiology*. 2017;8:1. doi:10.3389/fphys.2017.00001.
35. Choudhri AF, Siddiqui A, Khan NR, Cohen HL. Understanding bibliometric parameters and analysis. *Radiographics : a review publication of the Radiological Society of North America, Inc*. 2015;35(3):736-46. doi:10.1148/rg.2015140036.
36. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*. 2005;8(1):19-32. doi:10.1080/1364557032000119616.
37. Armstrong R, Hall BJ, Doyle J, Waters E. Cochrane Update. 'Scoping the scope' of a cochrane review. *Journal of public health*. 2011;33(1):147-50. doi:10.1093/pubmed/fdr015.
38. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implementation science : IS*. 2010;5:69. doi:10.1186/1748-5908-5-69.
39. Hortobágyi T, Granacher U, Fernandez-del-Olmo M. Whole body vibration and athletic performance: A scoping review. *European Journal of Human Movement*. 2014;33(2):1-25.
40. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther*. 2003;83(8):713-21.
41. Verhagen AP, de Vet HC, de Bie RA, Kessels AG, Boers M, Bouter LM et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *Journal of clinical epidemiology*. 1998;51(12):1235-41.
42. Schardt C, Adams MB, Owens T, Keitz S, Fontelo P. Utilization of the PICO framework to improve searching PubMed for clinical questions. *BMC medical informatics and decision making*. 2007;7:16. doi:10.1186/1472-6947-7-16.

43. Loturco I, Pereira LA, Kobal R, Maldonado T, Piazzzi AF, Bottino A et al. Improving sprint performance in soccer: effectiveness of jump squat and olympic push press exercises. *PLoS One*. 2016;11(4):e0153958. doi:10.1371/journal.pone.0153958.
44. Kobal R, Loturco I, Barroso R, Gil S, Cuniyochi R, Ugrinowitsch C et al. Effects of different combinations of strength, power, and plyometric training on the physical performance of elite young soccer players. *J Strength Cond Res*. 2017;31(6):1468-76. doi:10.1519/JSC.0000000000001609.
45. Yanci J, Castillo D, Iturricastillo A, Ayarra R, Nakamura FY. Effects of Two Different Volume-Equated Weekly Distributed Short-Term Plyometric Training Programs on Futsal Players' Physical Performance. *J Strength Cond Res*. 2016. doi:10.1519/JSC.0000000000001644.
46. Bruzas V, Kamandulis S, Venckunas T, Snieckus A, Mockus P. Effects of plyometric exercise training with external weights on punching ability of experienced amateur boxers. *J Sports Med Phys Fitness*. 2016.
47. Yanci J, Los Arcos A, Camara J, Castillo D, Garcia A, Castagna C. Effects of horizontal plyometric training volume on soccer players' performance. *Research in sports medicine (Print)*. 2016;24(4):308-19. doi:10.1080/15438627.2016.1222280.
48. Dello Iacono A, Martone D, Milic M, Padulo J. Vertical- vs. horizontal-oriented drop jump training: Chronic effects on explosive performances of elite handball players. *J Strength Cond Res*. 2017;31(4):921-31. doi:10.1519/JSC.0000000000001555.
49. Radnor JM, Lloyd RS, Oliver JL. Individual response to different forms of resistance training in school-aged boys. *J Strength Cond Res*. 2017;31(3):787-97. doi:10.1519/JSC.0000000000001527.
50. Kim YY, Min KO, Choi JH, Kim SH. The effects of sole vibration stimulation on Korean male professional volleyball players' jumping and balance ability. *J Phys Ther Sci*. 2016;28(5):1427-31. doi:10.1589/jpts.28.1427.
51. Asadi A, Ramirez-Campillo R, Meylan C, Nakamura FY, Canas-Jamet R, Izquierdo M. Effects of volume-based overload plyometric training on maximal-intensity exercise adaptations in young basketball players. *J Sports Med Phys Fitness*. 2016.
52. Hammami R, Granacher U, Makhlof I, Behm DG, Chaouachi A. Sequencing effects of balance and plyometric training on physical performance in youth soccer athletes. *J Strength Cond Res*. 2016;30(12):3278-89. doi:10.1519/JSC.0000000000001425.
53. Hammami M, Negra Y, Aouadi R, Shephard RJ, Chelly MS. Effects of an in-season plyometric training program on repeated change of direction and sprint performance in the junior soccer player. *J Strength Cond Res*. 2016;30(12):3312-20. doi:10.1519/JSC.0000000000001470.
54. Asadi A, Ramirez-Campillo R. Effects of cluster vs. traditional plyometric training sets on maximal-intensity exercise performance. *Medicina (Kaunas, Lithuania)*. 2016;52(1):41-5. doi:10.1016/j.medic.2016.01.001.
55. Weltin E, Gollhofer A, Mornieux G. Effects of perturbation or plyometric training on core control and knee joint loading in women during lateral movements. *Scand J Med Sci Sports*. 2017;27(3):299-308. doi:10.1111/sms.12657.

56. Hall E, Bishop DC, Gee TI. Effect of plyometric training on handspring vault performance and functional power in youth female gymnasts. *PLoS One*. 2016;11(2):e0148790. doi:10.1371/journal.pone.0148790.
57. de Hoyo M, Gonzalo-Skok O, Sanudo B, Carrascal C, Plaza-Armas JR, Camacho-Candil F et al. Comparative effects of in-season full-back squat, resisted sprint training, and plyometric training on explosive performance in U-19 elite soccer players. *J Strength Cond Res*. 2016;30(2):368-77. doi:10.1519/JSC.0000000000001094.
58. Speranza MJ, Gabbett TJ, Johnston RD, Sheppard JM. Effect of strength and power training on tackling ability in semiprofessional rugby league players. *J Strength Cond Res*. 2016;30(2):336-43. doi:10.1519/JSC.0000000000001058.
59. Chmielewski TL, George SZ, Tillman SM, Moser MW, Lentz TA, Indelicato PA et al. Low- versus high-intensity plyometric exercise during rehabilitation after anterior cruciate ligament reconstruction. *Am J Sports Med*. 2016;44(3):609-17. doi:10.1177/0363546515620583.
60. Ramírez-Campillo R, González-Jurado JA, Martínez C, Nakamura FY, Peñailillo L, Meylan CMP et al. Effects of plyometric training and creatine supplementation on maximal-intensity exercise and endurance in female soccer players. *Journal of Science and Medicine in Sport*. doi:10.1016/j.jsams.2015.10.005.
61. Nobre GG, Brito de Almeida M, Nobre IG, Karina Dos Santos F, Brinco RA, Arruda-Lima TR et al. Twelve-weeks of plyometric training improves motor performance of 7-10 year old overweight/obese boys: a randomized controlled intervention. *J Strength Cond Res*. 2016. doi:10.1519/JSC.0000000000001684.
62. Racil G, Zouhal H, Elmontassar W, Ben Abderrahmane A, De Sousa MV, Chamari K et al. Plyometric exercise combined with high-intensity interval training improves metabolic abnormalities in young obese females more so than interval training alone. *Appl Physiol Nutr Metab*. 2016;41(1):103-9. doi:10.1139/apnm-2015-0384.
63. Trecroci A, Cavaggioni L, Caccia R, Alberti G. Jump rope training: Balance and motor coordination in preadolescent soccer players. *J Sports Sci Med*. 2015;14(4):792-8.
64. Ache-Dias J, Dellagrana RA, Teixeira AS, Dal Pupo J, Moro AR. Effect of jumping interval training on neuromuscular and physiological parameters: a randomized controlled study. *Appl Physiol Nutr Metab*. 2016;41(1):20-5. doi:10.1139/apnm-2015-0368.
65. Lloyd RS, Radnor JM, De Ste Croix MB, Cronin JB, Oliver JL. Changes in sprint and jump performances after traditional, plyometric, and combined resistance training in male youth pre- and post-peak height velocity. *J Strength Cond Res*. 2016;30(5):1239-47. doi:10.1519/JSC.0000000000001216.
66. Jurado-Lavanant A, Alvero-Cruz JR, Pareja-Blanco F, Melero-Romero C, Rodriguez-Rosell D, Fernandez-Garcia JC. The effects of aquatic plyometric training on repeated jumps, drop jumps and muscle damage. *Int J Sports Med*. 2015. doi:10.1055/s-0034-1398574.

67. Loturco I, Pereira LA, Kobal R, Zanetti V, Kitamura K, Abad CC et al. Transference effect of vertical and horizontal plyometrics on sprint performance of high-level U-20 soccer players. *J Sports Sci.* 2015;33(20):2182-91. doi:10.1080/02640414.2015.1081394.
68. Pfile KR, Gribble PA, Buskirk GE, Meserth SM, Pietrosimone BG. Sustained improvements in dynamic balance and landing mechanics after a 6-week neuromuscular training program in college women's basketball players. *J Sport Rehabil.* 2016;25(3):233-40. doi:10.1123/jsr.2014-0323.
69. Pellegrino J, Ruby BC, Dumke CL. Effect of plyometrics on the energy cost of running and MHC and titin isoforms. *Med Sci Sports Exerc.* 2016;48(1):49-56. doi:10.1249/MSS.0000000000000747.
70. Fernandez-Fernandez J, Saez de Villarreal E, Sanz-Rivas D, Moya M. The effects of 8-week plyometric training on physical performance in young tennis players. *Pediatr Exerc Sci.* 2016;28(1):77-86. doi:10.1123/pes.2015-0019.
71. Rodriguez-Rosell D, Franco-Marquez F, Pareja-Blanco F, Mora-Custodio R, Yanez-Garcia JM, Gonzalez-Suarez JM et al. Effects of 6 weeks resistance training combined with plyometric and speed exercises on physical performance of pre-peak-height-velocity soccer players. *Int J Sports Physiol Perform.* 2016;11(2):240-6. doi:10.1123/ijsp.2015-0176.
72. Kijowski KN, Capps CR, Goodman CL, Erickson TM, Knorr DP, Triplett NT et al. Short-term resistance and plyometric training improves eccentric phase kinetics in jumping. *J Strength Cond Res.* 2015;29(8):2186-96. doi:10.1519/JSC.0000000000000904.
73. Ramirez-Campillo R, Vergara-Pedrerros M, Henriquez-Olguin C, Martinez-Salazar C, Alvarez C, Nakamura FY et al. Effects of plyometric training on maximal-intensity exercise and endurance in male and female soccer players. *J Sports Sci.* 2016;34(8):687-93. doi:10.1080/02640414.2015.1068439.
74. Franco-Marquez F, Rodriguez-Rosell D, Gonzalez-Suarez JM, Pareja-Blanco F, Mora-Custodio R, Yanez-Garcia JM et al. Effects of combined resistance training and plyometrics on physical performance in young soccer players. *Int J Sports Med.* 2015;36(11):906-14. doi:10.1055/s-0035-1548890.
75. Nyman E, Jr., Armstrong CW. Real-time feedback during drop landing training improves subsequent frontal and sagittal plane knee kinematics. *Clin Biomech (Bristol, Avon).* 2015;30(9):988-94. doi:10.1016/j.clinbiomech.2015.06.018.
76. McCormick BT, Hannon JC, Newton M, Shultz B, Detling N, Young WB. The effects of frontal- and sagittal-plane plyometrics on change-of-direction speed and power in adolescent female basketball players. *Int J Sports Physiol Perform.* 2016;11(1):102-7. doi:10.1123/ijsp.2015-0058.
77. Pereira A, Costa AM, Santos P, Figueiredo T, Joao PV. Training strategy of explosive strength in young female volleyball players. *Medicina (Kaunas, Lithuania).* 2015;51(2):126-31. doi:10.1016/j.medic.2015.03.004.

78. Tous-Fajardo J, Gonzalo-Skok O, Arjol-Serrano JL, Tesch P. Enhancing Change-of-Direction Speed in Soccer Players by Functional Inertial Eccentric Overload and Vibration Training. *Int J Sports Physiol Perform.* 2016;11(1):66-73. doi:10.1123/ijspp.2015-0010.
79. Carpentier A, Olbrechts N, Vieillevoys S, Poortmans JR. beta-Alanine supplementation slightly enhances repeated plyometric performance after high-intensity training in humans. *Amino Acids.* 2015;47(7):1479-83. doi:10.1007/s00726-015-1981-6.
80. Loturco I, Pereira LA, Kobal R, Zanetti V, Gil S, Kitamura K et al. Half-squat or jump squat training under optimum power load conditions to counteract power and speed decrements in Brazilian elite soccer players during the preseason. *J Sports Sci.* 2015;33(12):1283-92. doi:10.1080/02640414.2015.1022574.
81. Behrens M, Mau-Moeller A, Mueller K, Heise S, Gube M, Beuster N et al. Plyometric training improves voluntary activation and strength during isometric, concentric and eccentric contractions. *J Sci Med Sport.* 2016;19(2):170-6. doi:10.1016/j.jsams.2015.01.011.
82. Granacher U, Prieske O, Majewski M, Busch D, Muehlbauer T. The Role of Instability with Plyometric Training in Sub-elite Adolescent Soccer Players. *Int J Sports Med.* 2015;36(5):386-94. doi:10.1055/s-0034-1395519.
83. Chelly MS, Hermassi S, Shephard RJ. Effects of In-Season Short-term Plyometric Training Program on Sprint and Jump Performance of Young Male Track Athletes. *J Strength Cond Res.* 2015;29(8):2128-36. doi:10.1519/JSC.0000000000000860.
84. Saez de Villarreal E, Suarez-Arrones L, Requena B, Haff GG, Ferrete C. Effects of Plyometric and Sprint Training on Physical and Technical Skill Performance in Adolescent Soccer Players. *J Strength Cond Res.* 2015;29(7):1894-903. doi:10.1519/JSC.0000000000000838.
85. Mackala K, Fostiak M. Acute Effects of Plyometric Intervention-Performance Improvement and Related Changes in Sprinting Gait Variability. *J Strength Cond Res.* 2015;29(7):1956-65. doi:10.1519/JSC.0000000000000853.
86. Asadi A, Saez de Villarreal E, Arazi H. The effects of plyometric type neuromuscular training on postural control performance of male team basketball players. *J Strength Cond Res.* 2015;29(7):1870-5. doi:10.1519/JSC.0000000000000832.
87. Ramirez-Campillo R, Henriquez-Olguin C, Burgos C, Andrade DC, Zapata D, Martinez C et al. Effect of Progressive Volume-Based Overload During Plyometric Training on Explosive and Endurance Performance in Young Soccer Players. *J Strength Cond Res.* 2015;29(7):1884-93. doi:10.1519/JSC.0000000000000836.
88. Ramirez-Campillo R, Gallardo F, Henriquez-Olguin C, Meylan CM, Martinez C, Alvarez C et al. Effect of vertical, horizontal, and combined plyometric training on explosive, balance, and endurance performance of young soccer players. *J Strength Cond Res.* 2015;29(7):1784-95. doi:10.1519/JSC.0000000000000827.

89. Mendiguchia J, Martinez-Ruiz E, Morin JB, Samozino P, Edouard P, Alcaraz PE et al. Effects of hamstring-emphasized neuromuscular training on strength and sprinting mechanics in football players. *Scand J Med Sci Sports*. 2015;25(6):e621-9. doi:10.1111/sms.12388.
90. Ramachandran S, Pradhan B. Effects of short-term two weeks low intensity plyometrics combined with dynamic stretching training in improving vertical jump height and agility on trained basketball players. *Indian journal of physiology and pharmacology*. 2014;58(2):133-6.
91. Ramirez-Campillo R, Burgos CH, Henriquez-Olguin C, Andrade DC, Martinez C, Alvarez C et al. Effect of unilateral, bilateral, and combined plyometric training on explosive and endurance performance of young soccer players. *J Strength Cond Res*. 2015;29(5):1317-28. doi:10.1519/jsc.0000000000000762.
92. Saez de Villarreal E, Suarez-Arrones L, Requena B, Haff GG, Ramos Veliz R. Enhancing performance in professional water polo players: dryland training, in-water training, and combined training. *J Strength Cond Res*. 2015;29(4):1089-97. doi:10.1519/JSC.0000000000000707.
93. Arazi H, Mohammadi M, Asadi A. Muscular adaptations to depth jump plyometric training: Comparison of sand vs. land surface. *Interv Med Appl Sci*. 2014;6(3):125-30. doi:10.1556/IMAS.6.2014.3.5.
94. Carvalho A, Mourao P, Abade E. Effects of Strength Training Combined with Specific Plyometric exercises on body composition, vertical jump height and lower limb strength development in elite male handball players: a case study. *J Hum Kinet*. 2014;41:125-32. doi:10.2478/hukin-2014-0040.
95. Kibele A, Classen C, Muehlbauer T, Granacher U, Behm DG. Metastability in plyometric training on unstable surfaces: a pilot study. *BMC Sports Sci Med Rehabil*. 2014;6:30. doi:10.1186/2052-1847-6-30.
96. Russell RD, Nelson AG, Kraemer RR. Short bouts of high-intensity resistance-style training produce similar reductions in fasting blood glucose of diabetic offspring and controls. *J Strength Cond Res*. 2014;28(10):2760-7. doi:10.1519/JSC.0000000000000624.
97. Brito J, Vasconcellos F, Oliveira J, Krustup P, Rebelo A. Short-term performance effects of three different low-volume strength-training programmes in college male soccer players. *J Hum Kinet*. 2014;40:121-8. doi:10.2478/hukin-2014-0014.
98. Redondo JC, Alonso CJ, Sedano S, de Benito AM. Effects of a 12-week strength training program on experimented fencers' movement time. *J Strength Cond Res*. 2014. doi:10.1519/JSC.0000000000000581.
99. Attene G, Iuliano E, Di Cagno A, Calcagno G, Moalla W, Aquino G et al. Improving neuromuscular performance in young basketball players: plyometric vs. technique training. *J Sports Med Phys Fitness*. 2015;55(1-2):1-8.
100. Makaruk H, Czaplicki A, Sacewicz T, Sadowski J. The effects of single versus repeated plyometrics on landing biomechanics and jumping performance in men. *Biol Sport*. 2014;31(1):9-14. doi:10.5604/20831862.1083273.

101. Ozbar N, Ates S, Agopyan A. The effect of 8-week plyometric training on leg power, jump and sprint performance in female soccer players. *J Strength Cond Res.* 2014;28(10):2888-94. doi:10.1519/JSC.0000000000000541.
102. Ramirez-Campillo R, Andrade DC, Alvarez C, Henriquez-Olguin C, Martinez C, Baez-Sanmartin E et al. The effects of intersert rest on adaptation to 7 weeks of explosive training in young soccer players. *J Sports Sci Med.* 2014;13(2):287-96.
103. Lee C, Lee S, Yoo J. The effect of a complex training program on skating abilities in ice hockey players. *J Phys Ther Sci.* 2014;26(4):533-7. doi:10.1589/jpts.26.533.
104. Ramirez-Campillo R, Meylan C, Alvarez C, Henriquez-Olguin C, Martinez C, Canas-Jamett R et al. Effects of in-season low-volume high-intensity plyometric training on explosive actions and endurance of young soccer players. *J Strength Cond Res.* 2014;28(5):1335-42. doi:10.1519/jsc.0000000000000284.
105. Brown TN, Palmieri-Smith RM, McLean SG. Comparative adaptations of lower limb biomechanics during unilateral and bilateral landings after different neuromuscular-based ACL injury prevention protocols. *J Strength Cond Res.* 2014;28(10):2859-71. doi:10.1519/JSC.0000000000000472.
106. Garcia-Pinillos F, Martinez-Amat A, Hita-Contreras F, Martinez-Lopez EJ, Latorre-Roman PA. Effects of a contrast training program without external load on vertical jump, kicking speed, sprint, and agility of young soccer players. *J Strength Cond Res.* 2014;28(9):2452-60. doi:10.1519/JSC.0000000000000452.
107. Huang PY, Chen WL, Lin CF, Lee HJ. Lower extremity biomechanics in athletes with ankle instability after a 6-week integrated training program. *J Athl Train.* 2014;49(2):163-72. doi:10.4085/1062-6050-49.2.10.
108. Park GD, Lee JC, Lee J. The effect of low extremity plyometric training on back muscle power of high school throwing event athletes. *J Phys Ther Sci.* 2014;26(1):161-4. doi:10.1589/jpts.26.161.
109. Marques MC, Pereira A, Reis IG, van den Tillaar R. Does an in-season 6-week combined sprint and jump training program improve strength-speed abilities and kicking performance in young soccer players? *J Hum Kinet.* 2013;39:157-66. doi:10.2478/hukin-2013-0078
jhk-39-157 [pii].
110. Rebutini VZ, Pereira G, Bohrer RC, Ugrinowitsch C, Rodacki AL. Plyometric Long Jump Training With Progressive Loading Improves Kinetic and Kinematic Swimming Start Parameters. *J Strength Cond Res.* 2016;30(9):2392-8. doi:10.1519/JSC.0000000000000360.
111. Branislav R, Milivoj D, Abella CP, Deval VC, Sinisa K. Effects of combined and classic training on different isometric rate of force development parameters of leg extensors in female volleyball players: Discriminative analysis approach. *J Res Med Sci.* 2013;18(10):840-7.

112. Faigenbaum AD, Myer GD, Farrell A, Radler T, Fabiano M, Kang J et al. Integrative neuromuscular training and sex-specific fitness performance in 7-year-old children: an exploratory investigation. *J Athl Train.* 2014;49(2):145-53. doi:10.4085/1062-6050-49.1.08.
113. Sohnlein Q, Muller E, Stoggl TL. The effect of 16-week plyometric training on explosive actions in early to mid-puberty elite soccer players. *J Strength Cond Res.* 2014;28(8):2105-14. doi:10.1519/JSC.0000000000000387.
114. Stearns KM, Powers CM. Improvements in hip muscle performance result in increased use of the hip extensors and abductors during a landing task. *Am J Sports Med.* 2014;42(3):602-9. doi:10.1177/0363546513518410.
115. Macaluso F, Isaacs AW, Di Felice V, Myburgh KH. Acute change of titin at mid-sarcomere remains despite 8 wk of plyometric training. *J Appl Physiol (1985).* 2014;116(11):1512-9. doi:10.1152/jappphysiol.00420.2013.
116. Toivonen MH, Pollanen E, Ahtiainen M, Suominen H, Taaffe DR, Cheng S et al. OGT and OGA expression in postmenopausal skeletal muscle associates with hormone replacement therapy and muscle cross-sectional area. *Exp Gerontol.* 2013;48(12):1501-4. doi:10.1016/j.exger.2013.10.007.
117. Chaouachi A, Hammami R, Kaabi S, Chamari K, Drinkwater EJ, Behm DG. Olympic weightlifting and plyometric training with children provides similar or greater performance improvements than traditional resistance training. *J Strength Cond Res.* 2014;28(6):1483-96. doi:10.1519/JSC.0000000000000305.
118. Chelly MS, Hermassi S, Aouadi R, Shephard RJ. Effects of 8-week in-season plyometric training on upper and lower limb performance of elite adolescent handball players. *J Strength Cond Res.* 2014;28(5):1401-10. doi:10.1519/JSC.0000000000000279.
119. Lockie RG, Murphy AJ, Callaghan SJ, Jeffriess MD. Effects of sprint and plyometrics training on field sport acceleration technique. *J Strength Cond Res.* 2014;28(7):1790-801. doi:10.1519/JSC.0000000000000297.
120. Ramirez-Campillo R, Meylan CM, Alvarez-Lepin C, Henriquez-Olguin C, Martinez C, Andrade DC et al. The effects of interday rest on adaptation to 6 weeks of plyometric training in young soccer players. *J Strength Cond Res.* 2015;29(4):972-9. doi:10.1519/jsc.0000000000000283.
121. Baldon Rde M, Moreira Lobato DF, Yoshimatsu AP, dos Santos AF, Francisco AL, Pereira Santiago PR et al. Effect of plyometric training on lower limb biomechanics in females. *Clin J Sport Med.* 2014;24(1):44-50. doi:10.1097/01.jsm.0000432852.00391.de.
122. Marina M, Jemni M. Plyometric training performance in elite-oriented prepubertal female gymnasts. *J Strength Cond Res.* 2014;28(4):1015-25. doi:10.1519/JSC.0000000000000247.

123. Zribi A, Zouch M, Chaari H, Bouajina E, Ben Nasr H, Zaouali M et al. Short-term lower-body plyometric training improves whole body BMC, bone metabolic markers, and physical fitness in early pubertal male basketball players. *Pediatr Exerc Sci*. 2014;26(1):22-32. doi:10.1123/pes.2013-0053.
124. Behringer M, Neuerburg S, Matthews M, Mester J. Effects of two different resistance-training programs on mean tennis-serve velocity in adolescents. *Pediatr Exerc Sci*. 2013;25(3):370-84.
125. Behrens M, Mau-Moeller A, Bruhn S. Effect of plyometric training on neural and mechanical properties of the knee extensor muscles. *Int J Sports Med*. 2014;35(2):101-19. doi:10.1055/s-0033-1343401.
126. Ramirez-Campillo R, Alvarez C, Henriquez-Olguin C, Baez EB, Martinez C, Andrade DC et al. Effects of plyometric training on endurance and explosive strength performance in competitive middle- and long-distance runners. *J Strength Cond Res*. 2014;28(1):97-104. doi:10.1519/JSC.0b013e3182a1f44c.
127. Benito-Martinez E, Martinez-Amat A, Lara-Sanchez AJ, Berdejo-Del-Fresno D, Martinez-Lopez EJ. Effect of combined electrostimulation and plyometric training on 30 meters dash and triple jump. *J Sports Med Phys Fitness*. 2013;53(4):387-95.
128. Faude O, Roth R, Di Giovine D, Zahner L, Donath L. Combined strength and power training in high-level amateur football during the competitive season: a randomised-controlled trial. *J Sports Sci*. 2013;31(13):1460-7. doi:10.1080/02640414.2013.796065.
129. Pfile KR, Hart JM, Herman DC, Hertel J, Kerrigan DC, Ingersoll CD. Different exercise training interventions and drop-landing biomechanics in high school female athletes. *J Athl Train*. 2013;48(4):450-62. doi:10.4085/1062-6050-48.4.06.
130. Jafari M, Zolaktaf V, Marandi SM. Determination of the best pre-jump height for improvement of two-legged vertical jump. *International journal of preventive medicine*. 2013;4(Suppl 1):S104-9.
131. Vaczi M, Tollar J, Meszler B, Juhasz I, Karsai I. Short-term high intensity plyometric training program improves strength, power and agility in male soccer players. *J Hum Kinet*. 2013;36:17-26. doi:10.2478/hukin-2013-0002.
132. Barnes KR, Hopkins WG, McGuigan MR, Northuis ME, Kilding AE. Effects of resistance training on running economy and cross-country performance. *Med Sci Sports Exerc*. 2013;45(12):2322-31. doi:10.1249/MSS.0b013e31829af603.
133. Chaouachi A, Othman AB, Hammami R, Drinkwater EJ, Behm DG. The combination of plyometric and balance training improves sprint and shuttle run performances more often than plyometric-only training with children. *J Strength Cond Res*. 2014;28(2):401-12. doi:10.1519/JSC.0b013e3182987059.
134. Marshall BM, Moran KA. Which drop jump technique is most effective at enhancing countermovement jump ability, "countermovement" drop jump or "bounce" drop jump? *J Sports Sci*. 2013;31(12):1368-74. doi:10.1080/02640414.2013.789921.
135. Distefano LJ, Distefano MJ, Frank BS, Clark MA, Padua DA. Comparison of integrated and isolated training on performance measures and neuromuscular control. *J Strength Cond Res*. 2013;27(4):1083-90. doi:10.1519/JSC.0b013e318280d40b.

136. Leporace G, Praxedes J, Pereira GR, Pinto SM, Chagas D, Metsavaht L et al. Influence of a preventive training program on lower limb kinematics and vertical jump height of male volleyball athletes. *Phys Ther Sport*. 2013;14(1):35-43. doi:10.1016/j.ptsp.2012.02.005.
137. Sedano S, Marin PJ, Cuadrado G, Redondo JC. Concurrent training in elite male runners: the influence of strength versus muscular endurance training on performance outcomes. *J Strength Cond Res*. 2013;27(9):2433-43. doi:10.1519/JSC.0b013e318280cc26.
138. Ramirez-Campillo R, Andrade DC, Izquierdo M. Effects of plyometric training volume and training surface on explosive strength. *J Strength Cond Res*. 2013;27(10):2714-22. doi:10.1519/JSC.0b013e318280c9e9.
139. Houghton LA, Dawson BT, Rubenson J. Effects of plyometric training on achilles tendon properties and shuttle running during a simulated cricket batting innings. *J Strength Cond Res*. 2013;27(4):1036-46. doi:10.1519/JSC.0b013e3182651e7a.
140. Makaruk H, Porter JM, Czaplicki A, Sadowski J, Sacewicz T. The role of attentional focus in plyometric training. *J Sports Med Phys Fitness*. 2012;52(3):319-27.
141. Pienaar C, Coetzee B. Changes in selected physical, motor performance and anthropometric components of university-level rugby players after one microcycle of a combined rugby conditioning and plyometric training program. *J Strength Cond Res*. 2013;27(2):398-415. doi:10.1519/JSC.0b013e31825770ea.
142. Voelzke M, Stutzig N, Thorhauer HA, Granacher U. Promoting lower extremity strength in elite volleyball players: effects of two combined training methods. *J Sci Med Sport*. 2012;15(5):457-62. doi:10.1016/j.jsams.2012.02.004.
143. Cherif M, Said M, Chaatani S, Nejlaoui O, Gomri D, Abdallah A. The effect of a combined high-intensity plyometric and speed training program on the running and jumping ability of male handball players. *Asian J Sports Med*. 2012;3(1):21-8.
144. Michailidis Y, Fatouros IG, Primpa E, Michailidis C, Avloniti A, Chatzinikolaou A et al. Plyometrics' trainability in preadolescent soccer athletes. *J Strength Cond Res*. 2013;27(1):38-49. doi:10.1519/JSC.0b013e3182541ec6.
145. MacDonald CJ, Lamont HS, Garner JC. A comparison of the effects of 6 weeks of traditional resistance training, plyometric training, and complex training on measures of strength and anthropometrics. *J Strength Cond Res*. 2012;26(2):422-31. doi:10.1519/JSC.0b013e318220df79.
146. Kieffer HS, Lehman MA, Veacock D, Korkuch L. The effects of a short-term novel aquatic exercise program on functional strength and performance of older adults. *International journal of exercise science*. 2012;5(4):321-33.
147. Martinez-Lopez EJ, Benito-Martinez E, Hita-Contreras F, Lara-Sanchez A, Martinez-Amat A. Effects of electrostimulation and plyometric training program combination on jump height in teenage athletes. *J Sports Sci Med*. 2012;11(4):727-35.

148. Nakamura D, Suzuki T, Yasumatsu M, Akimoto T. Moderate running and plyometric training during off-season did not show a significant difference on soccer-related high-intensity performances compared with no-training controls. *J Strength Cond Res.* 2012;26(12):3392-7. doi:10.1519/JSC.0b013e3182474356.
149. Claudino JG, Mezencio B, Soncin R, Ferreira JC, Couto BP, Szmuchrowski LA. Pre vertical jump performance to regulate the training volume. *Int J Sports Med.* 2012;33(2):101-7. doi:10.1055/s-0031-1286293.
150. Kamandulis S, Snieckus A, Venckunas T, Aagaard P, Masiulis N, Skurvydas A. Rapid increase in training load affects markers of skeletal muscle damage and mechanical performance. *J Strength Cond Res.* 2012;26(11):2953-61. doi:10.1519/JSC.0b013e318243ff21.
151. Foure A, Nordez A, Cornu C. Effects of plyometric training on passive stiffness of gastrocnemii muscles and Achilles tendon. *Eur J Appl Physiol.* 2012;112(8):2849-57. doi:10.1007/s00421-011-2256-x.
152. Lockie RG, Murphy AJ, Scott BR, Janse de Jonge XA. Quantifying session ratings of perceived exertion for field-based speed training methods in team sport athletes. *J Strength Cond Res.* 2012;26(10):2721-8. doi:10.1519/JSC.0b013e3182429b0b.
153. Lloyd RS, Oliver JL, Hughes MG, Williams CA. The effects of 4-weeks of plyometric training on reactive strength index and leg stiffness in male youths. *J Strength Cond Res.* 2012;26(10):2812-9. doi:10.1519/JSC.0b013e318242d2ec.
154. Grieco CR, Cortes N, Greska EK, Lucci S, Onate JA. Effects of a combined resistance-plyometric training program on muscular strength, running economy, and Vo_{2peak} in division I female soccer players. *J Strength Cond Res.* 2012;26(9):2570-6. doi:10.1519/JSC.0b013e31823db1cf.
155. de Villarreal ES, Izquierdo M, Gonzalez-Badillo JJ. Enhancing jump performance after combined vs. maximal power, heavy-resistance, and plyometric training alone. *J Strength Cond Res.* 2011;25(12):3274-81. doi:10.1519/JSC.0b013e3182163085.
156. Makaruk H, Winchester JB, Sadowski J, Czaplicki A, Sacewicz T. Effects of unilateral and bilateral plyometric training on power and jumping ability in women. *J Strength Cond Res.* 2011;25(12):3311-8. doi:10.1519/JSC.0b013e318215fa33.
157. Greska EK, Cortes N, Van Lunen BL, Onate JA. A feedback inclusive neuromuscular training program alters frontal plane kinematics. *J Strength Cond Res.* 2012;26(6):1609-19. doi:10.1519/JSC.0b013e318234ebfb.
158. Lockie RG, Murphy AJ, Schultz AB, Knight TJ, Janse de Jonge XA. The effects of different speed training protocols on sprint acceleration kinematics and muscle strength and power in field sport athletes. *J Strength Cond Res.* 2012;26(6):1539-50. doi:10.1519/JSC.0b013e318234e8a0.
159. Alvarez M, Sedano S, Cuadrado G, Redondo JC. Effects of an 18-week strength training program on low-handicap golfers' performance. *J Strength Cond Res.* 2012;26(4):1110-21. doi:10.1519/JSC.0b013e31822dfa7d.
160. Tsang KK, DiPasquale AA. Improving the Q:H strength ratio in women using plyometric exercises. *J Strength Cond Res.* 2011;25(10):2740-5. doi:10.1519/JSC.0b013e31820d9e95.

161. Gonzalez-Aguero A, Vicente-Rodriguez G, Gomez-Cabello A, Ara I, Moreno LA, Casajus JA. A combined training intervention programme increases lean mass in youths with Down syndrome. *Research in developmental disabilities*. 2011;32(6):2383-8. doi:10.1016/j.ridd.2011.07.024.
162. Argus CK, Gill ND, Keogh JW, Blazevich AJ, Hopkins WG. Kinetic and training comparisons between assisted, resisted, and free countermovement jumps. *J Strength Cond Res*. 2011;25(8):2219-27. doi:10.1519/JSC.0b013e3181f6b0f4.
163. Taube W, Leukel C, Lauber B, Gollhofer A. The drop height determines neuromuscular adaptations and changes in jump performance in stretch-shortening cycle training. *Scand J Med Sci Sports*. 2012;22(5):671-83. doi:10.1111/j.1600-0838.2011.01293.x.
164. Sedano S, Matheu A, Redondo JC, Cuadrado G. Effects of plyometric training on explosive strength, acceleration capacity and kicking speed in young elite soccer players. *J Sports Med Phys Fitness*. 2011;51(1):50-8.
165. Roopchand-Martin S, Lue-Chin P. Plyometric training improves power and agility in Jamaica's national netball team. *West Indian Med J*. 2010;59(2):182-7.
166. Bonacci J, Green D, Saunders PU, Franettovich M, Blanch P, Vicenzino B. Plyometric training as an intervention to correct altered neuromotor control during running after cycling in triathletes: a preliminary randomised controlled trial. *Phys Ther Sport*. 2011;12(1):15-21. doi:10.1016/j.ptsp.2010.10.005.
167. Skurvydas A, Brazaitis M. Plyometric training does not affect central and peripheral muscle fatigue differently in prepubertal girls and boys. *Pediatr Exerc Sci*. 2010;22(4):547-56.
168. Potdevin FJ, Alberty ME, Chevutshi A, Pelayo P, Sidney MC. Effects of a 6-week plyometric training program on performances in pubescent swimmers. *J Strength Cond Res*. 2011;25(1):80-6. doi:10.1519/JSC.0b013e3181fef720.
169. Khelifa R, Aouadi R, Hermassi S, Chelly MS, Jlid MC, Hbacha H et al. Effects of a plyometric training program with and without added load on jumping ability in basketball players. *J Strength Cond Res*. 2010;24(11):2955-61. doi:10.1519/JSC.0b013e3181e37fbc.
170. Foue A, Nordez A, McNair P, Cornu C. Effects of plyometric training on both active and passive parts of the plantarflexors series elastic component stiffness of muscle-tendon complex. *Eur J Appl Physiol*. 2011;111(3):539-48. doi:10.1007/s00421-010-1667-4.
171. Chelly MS, Ghenem MA, Abid K, Hermassi S, Tabka Z, Shephard RJ. Effects of in-season short-term plyometric training program on leg power, jump- and sprint performance of soccer players. *J Strength Cond Res*. 2010;24(10):2670-6. doi:10.1519/JSC.0b013e3181e2728f.
172. Henry B, McLoda T, Docherty CL, Schrader J. The effect of plyometric training on peroneal latency. *J Sport Rehabil*. 2010;19(3):288-300.

173. Trzaskoma L, Tihanyi J, Trzaskoma Z. The effect of a short-term combined conditioning training for the development of leg strength and power. *J Strength Cond Res.* 2010;24(9):2498-505. doi:10.1519/JSC.0b013e3181e2e081.
174. Arabatzi F, Kellis E, Saez-Saez De Villarreal E. Vertical jump biomechanics after plyometric, weight lifting, and combined (weight lifting + plyometric) training. *J Strength Cond Res.* 2010;24(9):2440-8. doi:10.1519/JSC.0b013e3181e274ab.
175. Beaven CM, Gill ND, Ingram JR, Hopkins WG. Acute salivary hormone responses to complex exercise bouts. *J Strength Cond Res.* 2011;25(4):1072-8. doi:10.1519/JSC.0b013e3181bf4414.
176. Barber-Westin SD, Hermeto AA, Noyes FR. A six-week neuromuscular training program for competitive junior tennis players. *J Strength Cond Res.* 2010;24(9):2372-82. doi:10.1519/JSC.0b013e3181e8a47f.
177. King JA, Cipriani DJ. Comparing preseason frontal and sagittal plane plyometric programs on vertical jump height in high-school basketball players. *J Strength Cond Res.* 2010;24(8):2109-14. doi:10.1519/JSC.0b013e3181e347d1.
178. Saez Saez De Villarreal E, Requena B, Arampatzi F, Salonikidis K. Effect of plyometric training on chair-rise, jumping and sprinting performance in three age groups of women. *J Sports Med Phys Fitness.* 2010;50(2):166-73.
179. Foure A, Nordez A, Cornu C. Plyometric training effects on Achilles tendon stiffness and dissipative properties. *J Appl Physiol (1985).* 2010;109(3):849-54. doi:10.1152/jappphysiol.01150.2009.
180. Ismail MM, Ibrahim MM, Youssef EF, El Shorbagy KM. Plyometric training versus resistive exercises after acute lateral ankle sprain. *Foot Ankle Int.* 2010;31(6):523-30. doi:10.3113/FAI.2010.0523.
181. Berryman N, Maurel DB, Bosquet L. Effect of plyometric vs. dynamic weight training on the energy cost of running. *J Strength Cond Res.* 2010;24(7):1818-25. doi:10.1519/JSC.0b013e3181def1f5.
182. Herrero AJ, Martin J, Martin T, Abadia O, Fernandez B, Garcia-Lopez D. Short-term effect of plyometrics and strength training with and without superimposed electrical stimulation on muscle strength and anaerobic performance: A randomized controlled trial. Part II. *J Strength Cond Res.* 2010;24(6):1616-22. doi:10.1519/JSC.0b013e3181d8e84b.
183. Santos EJ, Janeira MA. The effects of plyometric training followed by detraining and reduced training periods on explosive strength in adolescent male basketball players. *J Strength Cond Res.* 2011;25(2):441-52. doi:10.1519/JSC.0b013e3181b62be3.
184. Skurvydas A, Brazaitis M, Streckis V, Rudas E. The effect of plyometric training on central and peripheral fatigue in boys. *Int J Sports Med.* 2010;31(7):451-7. doi:10.1055/s-0030-1251991.
185. Buchheit M, Mendez-Villanueva A, Delhomel G, Brughelli M, Ahmaidi S. Improving repeated sprint ability in young elite soccer players: repeated shuttle sprints vs. explosive strength training. *J Strength Cond Res.* 2010;24(10):2715-22. doi:10.1519/JSC.0b013e3181bf0223.
186. McLeod TC, Armstrong T, Miller M, Sauers JL. Balance improvements in female high school basketball players after a 6-week neuromuscular-training program. *J Sport Rehabil.* 2009;18(4):465-81.

187. Rubley MD, Haase AC, Holcomb WR, Girouard TJ, Tandy RD. The effect of plyometric training on power and kicking distance in female adolescent soccer players. *J Strength Cond Res.* 2011;25(1):129-34. doi:10.1519/JSC.0b013e3181b94a3d.
188. Ebben WP, Feldmann CR, Vanderzanden TL, Fauth ML, Petushek EJ. Periodized plyometric training is effective for women, and performance is not influenced by the length of post-training recovery. *J Strength Cond Res.* 2010;24(1):1-7. doi:10.1519/JSC.0b013e3181c49086.
189. Meylan C, Malatesta D. Effects of in-season plyometric training within soccer practice on explosive actions of young players. *J Strength Cond Res.* 2009;23(9):2605-13. doi:10.1519/JSC.0b013e3181b1f330.
190. Bishop DC, Smith RJ, Smith MF, Rigby HE. Effect of plyometric training on swimming block start performance in adolescents. *J Strength Cond Res.* 2009;23(7):2137-43. doi:10.1519/JSC.0b013e3181b866d0.
191. Sedano Campo S, Vaeyens R, Philippaerts RM, Redondo JC, de Benito AM, Cuadrado G. Effects of lower-limb plyometric training on body composition, explosive strength, and kicking speed in female soccer players. *J Strength Cond Res.* 2009;23(6):1714-22. doi:10.1519/JSC.0b013e3181b3f537.
192. Foure A, Nordez A, Guette M, Cornu C. Effects of plyometric training on passive stiffness of gastrocnemii and the musculo-articular complex of the ankle joint. *Scand J Med Sci Sports.* 2009;19(6):811-8. doi:10.1111/j.1600-0838.2008.00853.x.
193. Wu YK, Lien YH, Lin KH, Shih TT, Wang TG, Wang HK. Relationships between three potentiation effects of plyometric training and performance. *Scand J Med Sci Sports.* 2010;20(1):e80-6. doi:10.1111/j.1600-0838.2009.00908.x.
194. Carlson K, Magnusen M, Walters P. Effect of various training modalities on vertical jump. *Research in sports medicine (Print).* 2009;17(2):84-94. doi:10.1080/15438620902900351.
195. Farlinger CM, Fowles JR. The effect of sequence of skating-specific training on skating performance. *Int J Sports Physiol Perform.* 2008;3(2):185-98.
196. Guadalupe-Grau A, Perez-Gomez J, Olmedillas H, Chavarren J, Dorado C, Santana A et al. Strength training combined with plyometric jumps in adults: sex differences in fat-bone axis adaptations. *J Appl Physiol (1985).* 2009;106(4):1100-11. doi:10.1152/jappphysiol.91469.2008.
197. Potach DH, Katsavelis D, Karst GM, Latin RW, Stergiou N. The effects of a plyometric training program on the latency time of the quadriceps femoris and gastrocnemius short-latency responses. *J Sports Med Phys Fitness.* 2009;49(1):35-43.
198. Vescovi JD, Canavan PK, Hasson S. Effects of a plyometric program on vertical landing force and jumping performance in college women. *Phys Ther Sport.* 2008;9(4):185-92. doi:10.1016/j.ptsp.2008.08.001.
199. Thomas K, French D, Hayes PR. The effect of two plyometric training techniques on muscular power and agility in youth soccer players. *J Strength Cond Res.* 2009;23(1):332-5. doi:10.1519/JSC.0b013e318183a01a.

200. Vissing K, Brink M, Lonbro S, Sorensen H, Overgaard K, Danborg K et al. Muscle adaptations to plyometric vs. resistance training in untrained young men. *J Strength Cond Res.* 2008;22(6):1799-810. doi:10.1519/JSC.0b013e318185f673.
201. Grosset JF, Piscione J, Lambertz D, Perot C. Paired changes in electromechanical delay and musculo-tendinous stiffness after endurance or plyometric training. *Eur J Appl Physiol.* 2009;105(1):131-9. doi:10.1007/s00421-008-0882-8.
202. Marques MC, Tillaar R, Vescovi JD, Gonzalez-Badillo JJ. Changes in strength and power performance in elite senior female professional volleyball players during the in-season: a case study. *J Strength Cond Res.* 2008;22(4):1147-55. doi:10.1519/JSC.0b013e31816a42d0.
203. Perez-Gomez J, Olmedillas H, Delgado-Guerra S, Ara I, Vicente-Rodriguez G, Ortiz RA et al. Effects of weight lifting training combined with plyometric exercises on physical fitness, body composition, and knee extension velocity during kicking in football. *Appl Physiol Nutr Metab.* 2008;33(3):501-10. doi:10.1139/H08-026.
204. de Villarreal ES, Gonzalez-Badillo JJ, Izquierdo M. Low and moderate plyometric training frequency produces greater jumping and sprinting gains compared with high frequency. *J Strength Cond Res.* 2008;22(3):715-25. doi:10.1519/JSC.0b013e318163eade.
205. Rhea MR, Peterson MD, Lunt KT, Ayllon FN. The effectiveness of resisted jump training on the VertiMax in high school athletes. *J Strength Cond Res.* 2008;22(3):731-4. doi:10.1519/JSC.0b013e3181660c59.
206. Rhea MR, Peterson MD, Oliverson JR, Ayllon FN, Potenzianno BJ. An examination of training on the VertiMax resisted jumping device for improvements in lower body power in highly trained college athletes. *J Strength Cond Res.* 2008;22(3):735-40. doi:10.1519/JSC.0b013e3181660d61.
207. Ronnestad BR, Kvamme NH, Sunde A, Raastad T. Short-term effects of strength and plyometric training on sprint and jump performance in professional soccer players. *J Strength Cond Res.* 2008;22(3):773-80. doi:10.1519/JSC.0b013e31816a5e86.
208. Chappell JD, Limpisvasti O. Effect of a neuromuscular training program on the kinetics and kinematics of jumping tasks. *Am J Sports Med.* 2008;36(6):1081-6. doi:10.1177/0363546508314425.
209. Salonikidis K, Zafeiridis A. The effects of plyometric, tennis-drills, and combined training on reaction, lateral and linear speed, power, and strength in novice tennis players. *J Strength Cond Res.* 2008;22(1):182-91. doi:10.1519/JSC.0b013e31815f57ad.
210. Mangine GT, Ratamess NA, Hoffman JR, Faigenbaum AD, Kang J, Chilakos A. The effects of combined ballistic and heavy resistance training on maximal lower- and upper-body strength in recreationally trained men. *J Strength Cond Res.* 2008;22(1):132-9. doi:10.1519/JSC.0b013e31815f5729.
211. Mihalik JP, Libby JJ, Battaglini CL, McMurray RG. Comparing short-term complex and compound training programs on vertical jump height and power output. *J Strength Cond Res.* 2008;22(1):47-53. doi:10.1519/JSC.0b013e31815eee9e.

212. Dodd DJ, Alvar BA. Analysis of acute explosive training modalities to improve lower-body power in baseball players. *J Strength Cond Res.* 2007;21(4):1177-82. doi:10.1519/R-21306.1.
213. Kubo K, Morimoto M, Komuro T, Yata H, Tsunoda N, Kanehisa H et al. Effects of plyometric and weight training on muscle-tendon complex and jump performance. *Med Sci Sports Exerc.* 2007;39(10):1801-10. doi:10.1249/mss.0b013e31813e630a.
214. Ratamess NA, Kraemer WJ, Volek JS, French DN, Rubin MR, Gomez AL et al. The effects of ten weeks of resistance and combined plyometric/sprint training with the Meridian Elyte athletic shoe on muscular performance in women. *J Strength Cond Res.* 2007;21(3):882-7. doi:10.1519/R-50512.1.
215. Burgess KE, Connick MJ, Graham-Smith P, Pearson SJ. Plyometric vs. isometric training influences on tendon properties and muscle output. *J Strength Cond Res.* 2007;21(3):986-9. doi:10.1519/R-20235.1.
216. Stemm JD, Jacobson BH. Comparison of land- and aquatic-based plyometric training on vertical jump performance. *J Strength Cond Res.* 2007;21(2):568-71. doi:10.1519/R-20025.1.
217. Markovic G, Jukic I, Milanovic D, Metikos D. Effects of sprint and plyometric training on muscle function and athletic performance. *J Strength Cond Res.* 2007;21(2):543-9. doi:10.1519/R-19535.1.
218. Impellizzeri FM, Maffiuletti NA. Convergent evidence for construct validity of a 7-point likert scale of lower limb muscle soreness. *Clin J Sport Med.* 2007;17(6):494-6. doi:10.1097/JSM.0b013e31815aed57.
219. Faigenbaum AD, McFarland JE, Keiper FB, Tevlin W, Ratamess NA, Kang J et al. Effects of a short-term plyometric and resistance training program on fitness performance in boys age 12 to 15 years. *J Sports Sci Med.* 2007;6(4):519-25.
220. Saunders PU, Telford RD, Pyne DB, Peltola EM, Cunningham RB, Gore CJ et al. Short-term plyometric training improves running economy in highly trained middle and long distance runners. *J Strength Cond Res.* 2006;20(4):947-54. doi:10.1519/R-18235.1.
221. Malisoux L, Francaux M, Nielens H, Renard P, Lebacq J, Theisen D. Calcium sensitivity of human single muscle fibers following plyometric training. *Med Sci Sports Exerc.* 2006;38(11):1901-8. doi:10.1249/01.mss.0000232022.21361.47.
222. Pfeiffer RP, Shea KG, Roberts D, Grandstrand S, Bond L. Lack of effect of a knee ligament injury prevention program on the incidence of noncontact anterior cruciate ligament injury. *J Bone Joint Surg Am.* 2006;88(8):1769-74. doi:10.2106/JBJS.E.00616.
223. Herrero JA, Izquierdo M, Maffiuletti NA, Garcia-Lopez J. Electromyostimulation and plyometric training effects on jumping and sprint time. *Int J Sports Med.* 2006;27(7):533-9. doi:10.1055/s-2005-865845.
224. Kotzamanidis C. Effect of plyometric training on running performance and vertical jumping in prepubertal boys. *J Strength Cond Res.* 2006;20(2):441-5. doi:10.1519/R-16194.1.

225. Miller MG, Herniman JJ, Ricard MD, Cheatham CC, Michael TJ. The effects of a 6-week plyometric training program on agility. *J Sport Sci Med*. 2006;5(3):459-65.
226. Diracoglu D, Aydin R, Baskent A, Celik A. Effects of kinesthesia and balance exercises in knee osteoarthritis. *J Clin Rheumatol*. 2005;11(6):303-10.
227. Lephart SM, Abt JP, Ferris CM, Sell TC, Nagai T, Myers JB et al. Neuromuscular and biomechanical characteristic changes in high school athletes: a plyometric versus basic resistance program. *Br J Sports Med*. 2005;39(12):932-8. doi:39/12/932 [pii] 10.1136/bjism.2005.019083.
228. Moore EW, Hickey MS, Reiser RF. Comparison of two twelve week off-season combined training programs on entry level collegiate soccer players' performance. *J Strength Cond Res*. 2005;19(4):791-8. doi:10.1519/R-15384.1.
229. Myer GD, Ford KR, Brent JL, Hewett TE. The effects of plyometric vs. dynamic stabilization and balance training on power, balance, and landing force in female athletes. *J Strength Cond Res*. 2006;20(2):345-53. doi:10.1519/R-17955.1.
230. Martel GF, Harmer ML, Logan JM, Parker CB. Aquatic plyometric training increases vertical jump in female volleyball players. *Med Sci Sports Exerc*. 2005;37(10):1814-9.
231. McCurdy KW, Langford GA, Doscher MW, Wiley LP, Mallard KG. The effects of short-term unilateral and bilateral lower-body resistance training on measures of strength and power. *J Strength Cond Res*. 2005;19(1):9-15. doi:10.1519/14173.1.
232. Myer GD, Ford KR, Palumbo JP, Hewett TE. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J Strength Cond Res*. 2005;19(1):51-60. doi:10.1519/13643.1.
233. Irmischer BS, Harris C, Pfeiffer RP, DeBeliso MA, Adams KJ, Shea KG. Effects of a knee ligament injury prevention exercise program on impact forces in women. *J Strength Cond Res*. 2004;18(4):703-7. doi:10.1519/R-13473.1.
234. Canavan PK, Vescovi JD. Evaluation of power prediction equations: peak vertical jumping power in women. *Med Sci Sports Exerc*. 2004;36(9):1589-93.
235. Chimera NJ, Swanik KA, Swanik CB, Straub SJ. Effects of Plyometric Training on Muscle-Activation Strategies and Performance in Female Athletes. *J Athl Train*. 2004;39(1):24-31.
236. Wilkerson GB, Colston MA, Short NI, Neal KL, Hoewischer PE, Pixley JJ. Neuromuscular Changes in Female Collegiate Athletes Resulting From a Plyometric Jump-Training Program. *J Athl Train*. 2004;39(1):17-23.
237. Robinson LE, Devor ST, Merrick MA, Buckworth J. The effects of land vs. aquatic plyometrics on power, torque, velocity, and muscle soreness in women. *J Strength Cond Res*. 2004;18(1):84-91.
238. Luebbers PE, Potteiger JA, Hulver MW, Thyfault JP, Carper MJ, Lockwood RH. Effects of plyometric training and recovery on vertical jump performance and anaerobic power. *J Strength Cond Res*. 2003;17(4):704-9.

239. Buckley JD, Brinkworth GD, Abbott MJ. Effect of bovine colostrum on anaerobic exercise performance and plasma insulin-like growth factor I. *J Sports Sci.* 2003;21(7):577-88. doi:10.1080/0264041031000101935.
240. Siegler J, Gaskill S, Ruby B. Changes evaluated in soccer-specific power endurance either with or without a 10-week, in-season, intermittent, high-intensity training protocol. *J Strength Cond Res.* 2003;17(2):379-87.
241. Spurrs RW, Murphy AJ, Watsford ML. The effect of plyometric training on distance running performance. *Eur J Appl Physiol.* 2003;89(1):1-7. doi:10.1007/s00421-002-0741-y.
242. Turner AM, Owings M, Schwane JA. Improvement in running economy after 6 weeks of plyometric training. *J Strength Cond Res.* 2003;17(1):60-7.
243. Maffiuletti NA, Dugnani S, Folz M, Di Pierno E, Mauro F. Effect of combined electrostimulation and plyometric training on vertical jump height. *Med Sci Sports Exerc.* 2002;34(10):1638-44. doi:10.1249/01.MSS.0000031481.28915.56.
244. McLaughlin EJ. A comparison between two training programs and their effects on fatigue rates in women. *J Strength Cond Res.* 2001;15(1):25-9.
245. Diallo O, Dore E, Duche P, Van Praagh E. Effects of plyometric training followed by a reduced training programme on physical performance in prepubescent soccer players. *J Sports Med Phys Fitness.* 2001;41(3):342-8.
246. Matavulj D, Kukolj M, Ugarkovic D, Tihanyi J, Jaric S. Effects of plyometric training on jumping performance in junior basketball players. *J Sports Med Phys Fitness.* 2001;41(2):159-64.
247. Kraemer WJ, Mazzetti SA, Nindl BC, Gotshalk LA, Volek JS, Bush JA et al. Effect of resistance training on women's strength/power and occupational performances. *Med Sci Sports Exerc.* 2001;33(6):1011-25.
248. Witzke KA, Snow CM. Effects of plyometric jump training on bone mass in adolescent girls. *Med Sci Sports Exerc.* 2000;32(6):1051-7.
249. Hewett TE, Lindenfeld TN, Riccobene JV, Noyes FR. The effect of neuromuscular training on the incidence of knee injury in female athletes. A prospective study. *Am J Sports Med.* 1999;27(6):699-706. doi:10.1177/03635465990270060301.
250. Cossor JM, Blanksby BA, Elliott BC. The influence of plyometric training on the freestyle tumble turn. *J Sci Med Sport.* 1999;2(2):106-16.
251. Newton RU, Kraemer WJ, Hakkinen K. Effects of ballistic training on preseason preparation of elite volleyball players. *Med Sci Sports Exerc.* 1999;31(2):323-30.
252. Cornu C, Almeida Silveira MI, Goubel F. Influence of plyometric training on the mechanical impedance of the human ankle joint. *Eur J Appl Physiol Occup Physiol.* 1997;76(3):282-8.
253. Hewett TE, Stroupe AL, Nance TA, Noyes FR. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996;24(6):765-73. doi:10.1177/036354659602400611.

254. Wilson GJ, Murphy AJ, Giorgi A. Weight and plyometric training: effects on eccentric and concentric force production. *Can J Appl Physiol.* 1996;21(4):301-15.
255. Wilson GJ, Murphy AJ. Strength diagnosis: the use of test data to determine specific strength training. *J Sports Sci.* 1996;14(2):167-73. doi:10.1080/02640419608727698.
256. Fowler NE, Trzaskoma Z, Wit A, Iskra L, Lees A. The effectiveness of a pendulum swing for the development of leg strength and counter-movement jump performance. *J Sports Sci.* 1995;13(2):101-8. doi:10.1080/02640419508732217.
257. Kramer JF, Morrow A, Leger A. Changes in rowing ergometer, weight lifting, vertical jump and isokinetic performance in response to standard and standard plus plyometric training programs. *Int J Sports Med.* 1993;14(8):449-54. doi:10.1055/s-2007-1021209.
258. Wilson GJ, Newton RU, Murphy AJ, Humphries BJ. The optimal training load for the development of dynamic athletic performance. *Med Sci Sports Exerc.* 1993;25(11):1279-86.
259. Brown ME, Mayhew JL, Boleach LW. Effect of plyometric training on vertical jump performance in high school basketball players. *J Sports Med Phys Fitness.* 1986;26(1):1-4.
260. Ford HT, Jr., Puckett JR, Drummond JP, Sawyer K, Gantt K, Fussell C. Effects of three combinations of plyometric and weight training programs on selected physical fitness test items. *Percept Mot Skills.* 1983;56(3):919-22. doi:10.2466/pms.1983.56.3.919.
261. Bogdanis GC, Tsoukos A, Kaloheri O, Terzis G, Veligeas P, Brown LE. Comparison between Unilateral and Bilateral Plyometric Training on Single and Double Leg Jumping Performance and Strength. *J Strength Cond Res.* 2017. doi:10.1519/JSC.0000000000001962.
262. Giovanelli N, Taboga P, Rejc E, Lazzar S. Effects of strength, explosive and plyometric training on energy cost of running in ultra-endurance athletes. *Eur J Sport Sci.* 2017;1-9. doi:10.1080/17461391.2017.1305454.
263. Jeffreys M, De Ste Croix M, Lloyd RS, Oliver JL, Hughes J. The effect of varying plyometric volume on stretch-shortening cycle capability in collegiate male rugby players. *J Strength Cond Res.* 2017. doi:10.1519/JSC.0000000000001907.
264. Chaouachi M, Granacher U, Makhlof I, Hammami R, Behm DG, Chaouachi A. Within Session Sequence of Balance and Plyometric Exercises Does Not Affect Training Adaptations with Youth Soccer Athletes. *J Sports Sci Med.* 2017;16(1):125-36.
265. Negra Y, Chaabene H, Sammoud S, Bouguezzi R, Abbes MA, Hachana Y et al. Effects of Plyometric Training on Physical Fitness in Prepuberal Soccer Athletes. *Int J Sports Med.* 2017;38(5):370-7. doi:10.1055/s-0042-122337.
266. Assuncao AR, Bottaro M, Cardoso EA, D PDdS, Ferraz M, Vieira CA et al. Effects of a low-volume plyometric training in anaerobic performance of adolescent athletes. *J Sports Med Phys Fitness.* 2017. doi:10.23736/S0022-4707.17.07173-0.

267. Chaabene H, Negra Y. The Effect of Plyometric Training Volume in Prepubertal Male Soccer Players' Athletic Performance. *Int J Sports Physiol Perform*. 2017;1-22. doi:10.1123/ijsp.2016-0372.
268. Hirayama K, Iwanuma S, Ikeda N, Yoshikawa A, Ema R, Kawakami Y. Plyometric Training Favors Optimizing Muscle-Tendon Behavior during Depth Jumping. *Frontiers in physiology*. 2017;8:16. doi:10.3389/fphys.2017.00016.
269. Rodriguez-Rosell D, Torres-Torrel J, Franco-Marquez F, Gonzalez-Suarez JM, Gonzalez-Badillo JJ. Effects of light-load maximal lifting velocity weight training vs. combined weight training and plyometrics on sprint, vertical jump and strength performance in adult soccer players. *J Sci Med Sport*. 2017. doi:10.1016/j.jsams.2016.11.010.
270. Egan-Shuttler JD, Edmonds R, Eddy C, O'Neill V, Ives SJ. The Effect of Concurrent Plyometric Training Versus Submaximal Aerobic Cycling on Rowing Economy, Peak Power, and Performance in Male High School Rowers. *Sports medicine - open*. 2017;3(1):7. doi:10.1186/s40798-017-0075-2.
271. Daehlin TE, Haugen OC, Haugerud S, Hollan I, Raastad T, Ronnestad BR. Combined Plyometric & Strength Training Improves Ice-hockey Players` On-ice Sprint. *Int J Sports Physiol Perform*. 2016:1-22. doi:10.1123/ijsp.2016-0262.
272. Moran J, Sandercock GRH, Ramirez-Campillo R, Todd O, Collison J, Parry DA. Maturation-Related Effect of Low-Dose Plyometric Training on Performance in Youth Hockey Players. *Pediatr Exerc Sci*. 2017;29(2):194-202. doi:10.1123/pes.2016-0151.
273. Benis R, Bonato M, La Torre A. Elite Female Basketball Players' Body-Weight Neuromuscular Training and Performance on the Y-Balance Test. *J Athl Train*. 2016;51(9):688-95. doi:10.4085/1062-6050-51.12.03.
274. Khodaei K, Mohammadi A, Badri N. A comparison of assisted, resisted, and common plyometric training modes to enhance sprint and agility performance. *J Sports Med Phys Fitness*. 2017. doi:10.23736/S0022-4707.17.06901-8.
275. Arabatzi F. Adaptations in movement performance after plyometric training on mini-trampoline in children. *J Sports Med Phys Fitness*. 2016.
276. Stark L, Pickett K, Bird M, King AC. Influence of Knee-to-Feet Jump Training on Vertical Jump and Hang Clean Performance. *J Strength Cond Res*. 2016;30(11):3084-9. doi:10.1519/JSC.0000000000001403.
277. Teo SY, Newton MJ, Newton RU, Dempsey AR, Fairchild TJ. Comparing the Effectiveness of a Short-Term Vertical Jump vs. Weightlifting Program on Athletic Power Development. *J Strength Cond Res*. 2016;30(10):2741-8. doi:10.1519/JSC.0000000000001379.
278. Kim YY, Park SE. Comparison of whole-body vibration exercise and plyometric exercise to improve isokinetic muscular strength, jumping performance and balance of female volleyball players. *J Phys Ther Sci*. 2016;28(11):3140-4. doi:10.1589/jpts.28.3140.

279. Myer GD, Ford KR, McLean SG, Hewett TE. The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics. *Am J Sports Med.* 2006;34(3):445-55. doi:10.1177/0363546505281241.
280. Impellizzeri FM, Rampinini E, Castagna C, Martino F, Fiorini S, Wisloff U. Effect of plyometric training on sand versus grass on muscle soreness and jumping and sprinting ability in soccer players. *Br J Sports Med.* 2008;42(1):42-6. doi:10.1136/bjism.2007.038497.
281. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3-13. doi:10.1249/MSS.0b013e31818cb278.
282. Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. *Sportscience.* 2005;9:6-13.
283. Greenland S, Senn SJ, Rothman KJ, Carlin JB, Poole C, Goodman SN et al. Statistical tests, P values, confidence intervals, and power: a guide to misinterpretations. *Eur J Epidemiol.* 2016;31(4):337-50. doi:10.1007/s10654-016-0149-3.
284. Rhea MR. Determining the magnitude of treatment effects in strength training research through the use of the effect size. *J Strength Cond Res.* 2004;18(4):918-20. doi:10.1519/14403.1.
285. Datson N, Hulton A, Andersson H, Lewis T, Weston M, Drust B et al. Applied physiology of female soccer: an update. *Sports Med.* 2014;44(9):1225-40. doi:10.1007/s40279-014-0199-1.
286. Legerlotz K, Marzilger R, Bohm S, Arampatzis A. Physiological adaptations following resistance training in youth athletes- a narrative review. *Pediatr Exerc Sci.* 2016;28(4):501-20. doi:10.1123/pes.2016-0023.
287. Mersmann F, Bohm S, Schroll A, Boeth H, Duda G, Arampatzis A. Evidence of imbalanced adaptation between muscle and tendon in adolescent athletes. *Scand J Med Sci Sports.* 2014;24(4):e283-9. doi:10.1111/sms.12166.
288. Mersmann F, Bohm S, Schroll A, Boeth H, Duda GN, Arampatzis A. Muscle and tendon adaptation in adolescent athletes: A longitudinal study. *Scand J Med Sci Sports.* 2017;27(1):75-82. doi:10.1111/sms.12631.
289. Ebben WP. Complex training: a brief review. *J Sports Sci Med.* 2002;1(2):42-6.
290. Ebben WP, Blackard DO. Strength and conditioning practices of National Football League strength and conditioning coaches. *J Strength Cond Res.* 2001;15(1):48-58.
291. Ebben WP, Carroll RM, Simenz CJ. Strength and conditioning practices of National Hockey League strength and conditioning coaches. *J Strength Cond Res.* 2004;18(4):889-97. doi:10.1519/14133.1.
292. Ebben WP, Hintz MJ, Simenz CJ. Strength and conditioning practices of Major League Baseball strength and conditioning coaches. *J Strength Cond Res.* 2005;19(3):538-46. doi:10.1519/R-15464.1.
293. Ebben WP, Watts PB. A review of combined weight training and plyometric training modes: Complex training. *Strength Cond J.* 1998;20(5):18-27.

294. Winwood PW, Keogh JW, Harris NK. The strength and conditioning practices of strongman competitors. *J Strength Cond Res.* 2011;25(11):3118-28. doi:10.1519/JSC.0b013e318212daea.
295. Swinton PA, Lloyd R, Agouris I, Stewart A. Contemporary training practices in elite British powerlifters: survey results from an international competition. *J Strength Cond Res.* 2009;23(2):380-4. doi:10.1519/JSC.0b013e31819424bd.
296. Simenz CJ, Dugan CA, Ebben WP. Strength and conditioning practices of National Basketball Association strength and conditioning coaches. *J Strength Cond Res.* 2005;19(3):495-504. doi:10.1519/15264.1.
297. Ziv G, Lidor R. Vertical jump in female and male volleyball players: a review of observational and experimental studies. *Scand J Med Sci Sports.* 2010;20(4):556-67. doi:10.1111/j.1600-0838.2009.01083.x.
298. Rossler R, Donath L, Verhagen E, Junge A, Schweizer T, Faude O. Exercise-based injury prevention in child and adolescent sport: a systematic review and meta-analysis. *Sports Med.* 2014;44(12):1733-48. doi:10.1007/s40279-014-0234-2.
299. Bobbert MF. Drop jumping as a training method for jumping ability. *Sports Med.* 1990;9(1):7-22.
300. Giatsis G, Kollias I, Panoutsakopoulos V, Papaiaikovou G. Biomechanical differences in elite beach-volleyball players in vertical squat jump on rigid and sand surface. *Sports biomechanics.* 2004;3(1):145-58. doi:10.1080/14763140408522835.
301. Büsch D, Pabst J, Mühlbauer T, Ehrhardt P, Granacher U. Effects of plyometric training using unstable surfaces on jump and sprint performance in young sub-elite handball players. *Sports Orthop Traumatol.* 2015;31(4):299-308.
302. Negra Y, Chaabene H, Stöggl T, Hammami M, Chelly MS, Hachana Y. Effectiveness and time-course adaptation of resistance training vs. plyometric training in prepubertal soccer players. *Journal of Sport and Health Science.* IN PRESS. doi:10.1016/j.jshs.2016.07.008.
303. Visnes H, Bahr R. Training volume and body composition as risk factors for developing jumper's knee among young elite volleyball players. *Scand J Med Sci Sports.* 2013;23(5):607-13. doi:10.1111/j.1600-0838.2011.01430.x.
304. Lesinski M, Prieske O, Granacher U. Effects and dose-response relationships of resistance training on physical performance in youth athletes: a systematic review and meta-analysis. *Br J Sports Med.* 2016;50(13):781-95. doi:10.1136/bjsports-2015-095497.
305. Ebben WP. Practical guidelines for plyometric intensity. *NSCA's Performance Training Journal.* 2007;6:12-6.
306. Peng HT, Kernozek TW, Song CY. Quadriceps and hamstring activation during drop jumps with changes in drop height. *Phys Ther Sport.* 2011;12(3):127-32. doi:10.1016/j.ptsp.2010.10.001.
307. Chu D. *Jumping into plyometrics.* Human Kinetics. 1998;Champaign:IL.
308. Amara S, Mkaouer B, Chaabene H, Negra Y, Hammami M, Bouguezzi R. Effect of plyometric training on young athlete's performance according to body corpulence and dropping height. *J Athl Enhancement.* 2015;4(4). doi:10.4172/2324-9080.1000203.

309. Bobbert MF, Huijing PA, van Ingen Schenau GJ. Drop jumping. II. The influence of dropping height on the biomechanics of drop jumping. *Med Sci Sports Exerc.* 1987;19(4):339-46.
310. Bobbert MF, Huijing PA, van Ingen Schenau GJ. Drop jumping. I. The influence of jumping technique on the biomechanics of jumping. *Med Sci Sports Exerc.* 1987;19(4):332-8.
311. Ebben WP, Simenz C, Jensen RL. Evaluation of plyometric intensity using electromyography. *J Strength Cond Res.* 2008;22(3):861-8. doi:10.1519/JSC.0b013e31816a834b.
312. Ruan M, Li L. Approach run increases preactivation and eccentric phases muscle activity during drop jumps from different drop heights. *J Electromyogr Kinesiol.* 2010;20(5):932-8. doi:10.1016/j.jelekin.2009.08.007.
313. Andrade DC, Manzo O, Beltrán AR, Alvares C, Del Río R, Toledo C et al. Kinematic and neuromuscular measures of intensity during plyometric jumps. *Journal of Strength and Conditioning Research.* IN PRESS.
314. Flanagan EP, Comyns TM. The use of contact time and the reactive strength index to optimize fast stretch-shortening cycle training. *Strength & Conditioning Journal.* 2008;30(5):32-8 doi:10.1519/SSC.0b013e318187e25b.
315. Kenney WL, Wilmore J, Costill D. *Physiology of Sport and Exercise.* Human Kinetics, Champaign, IL. 2015.
316. Houmard JA, Johns RA. Effects of taper on swim performance. Practical implications. *Sports Med.* 1994;17(4):224-32.
317. Mujika I, Padilla S, Pyne D, Busso T. Physiological changes associated with the pre-event taper in athletes. *Sports Med.* 2004;34(13):891-927.
318. Neary JP, Martin TP, Quinney HA. Effects of taper on endurance cycling capacity and single muscle fiber properties. *Med Sci Sports Exerc.* 2003;35(11):1875-81. doi:10.1249/01.MSS.0000093617.28237.20.
319. Read MM, Cisar C. The influence of varied rest interval lengths on depth jump performance. *J Strength Cond Res.* 2001;15(3):279-83.
320. Kraemer WJ, Ratamess NA. Hormonal responses and adaptations to resistance exercise and training. *Sports Med.* 2005;35(4):339-61. doi:3544 [pii].
321. Balsom PD, Seger JY, Sjodin B, Ekblom B. Maximal-intensity intermittent exercise: effect of recovery duration. *Int J Sports Med.* 1992;13(7):528-33. doi:10.1055/s-2007-1021311.
322. Balsom PD, Seger JY, Sjodin B, Ekblom B. Physiological responses to maximal intensity intermittent exercise. *Eur J Appl Physiol Occup Physiol.* 1992;65(2):144-9.
323. Marginson V, Rowlands AV, Gleeson NP, Eston RG. Comparison of the symptoms of exercise-induced muscle damage after an initial and repeated bout of plyometric exercise in men and boys. *J Appl Physiol.* 2005;99(3):1174-81. doi:01193.2004 [pii]
10.1152/jappphysiol.01193.2004.

324. Wathen D. Literature Review: Explosive/Plyometric Exercises. *Strength & Conditioning Journal*. 1993;15(3):17-9.