

The influence of maturation on the reliability of the Nordic hamstring exercise in male youth footballers

Fernandes, John; Moran, Jason; Clarke, Hannah; Drury, Ben

Published in:
Translational Sports Medicine

Publication date:
2020

The re-use license for this item is:
CC BY-NC-ND

This document version is the:
Peer reviewed version

The final published version is available direct from the publisher website at:
[10.1002/tsm2.124](https://doi.org/10.1002/tsm2.124)

[Find this output at Hartpury Pure](#)

Citation for published version (APA):
Fernandes, J., Moran, J., Clarke, H., & Drury, B. (2020). The influence of maturation on the reliability of the Nordic hamstring exercise in male youth footballers. *Translational Sports Medicine*, 3(2), 148-153.
<https://doi.org/10.1002/tsm2.124>

1 **TRANSLATIONAL SPORTS MEDICINE**

2
3 **Title:** The influence of maturation on the reliability of the Nordic hamstring exercise in male
4 youth footballers

5
6 **Running head:** Maturation and NHE reliability

7
8 **Authors:** John F. T. Fernandes¹, Jason Moran², Hannah Clarke¹, Benjamin Drury¹

9
10 **Author affiliations:** ¹Higher Education Sport, Hartpury University, Hartpury, UK.

11 ²School of Sport, Rehabilitation, and Exercise Sciences, University of Essex, Colchester, UK

12
13 **Corresponding author:** John Fernandes, Sport, Health and Well-being Arena, Hartpury
14 University, Hartpury, GL19 3BE. 01452 702269. jfmtfernandes@hotmail.co.uk.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Abstract

This study sought to establish the reliability of the Nordic hamstring exercise (NHE) in male youth football players. Sixty-four youth football players completed two x three repetitions of the NHE, separated by one week. Eccentric hamstring strength was measured during the NHE using the NordBord. Participants were categorised via maturity offset (based on peak height velocity [PHV]) and age. For all dependent variables and groups, the typical error (TE) was greater than the smallest worthwhile change. Reliability for left, right, bilateral and relative peak force for the U11s (TE=0.26 to 11.1N, coefficient of variation (CV) = 5.9 to 7.4%), U13s (TE=0.28 to 17.9N, CVs=5.6 to 7.8%) and U16s (TE=0.28 to 24.3, CVs=6.6 to 8.7%) was favourable and demonstrated no clear pattern between groups. According to PHV, those less mature provided smaller TEs (0.22 to 9.3N) and CVs (4.8 to 5.7%) compared to their more mature counterparts (TE=0.30 to 22.5N, CVs=7.2 to 8.5%). For all age and maturation groups, imbalances yielded poor reliability (TE=7.1 to 10.8N, CVs=33.1 to 38.3%). Eccentric left and right limb, bilateral and relative hamstring peak force can reliably be measured during the NHE across maturation stages. Applied practitioners should exercise caution when assessing muscular imbalances using the NHE.

Key words

Peak force; imbalance; reproducibility; relative force; eccentric; eccentric force

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Introduction

Strength refers to the maximal force or torque that the skeletal muscles can exert¹ and is dependent on the muscle contraction type (e.g. concentric, isometric or eccentric) and velocity. For example, when compared to eccentric contractions, concentric and isometric contractions exhibit lower peak forces². Furthermore, for the youth athlete, strength is an important part of sporting performance³. That is, strength is moderately and strongly correlated with sprint ($r = > -0.60$) and jump performance ($r = > -0.76$), respectively⁴ and is integral for the performance of fundamental movement skills³.

Strength increases naturally throughout maturation⁵ and is underpinned by neural and muscular alterations⁶⁻⁹. For those who want to maximise a youth's strength gain, resistance training is a potent method for enhancing strength¹⁰. However, the magnitude of strength adaptations to resistance training can be dependent on maturation status, amongst other factors¹⁰. For example, strength adaptations are greater in boys during, and after, peak height velocity (PHV) than before PHV¹¹. When resistance training (e.g. traditional high intensity training, Olympic lifting) youths in the pre-, mid- and post-PHV stages can expect enhancements in strength that range from 3.5 to 36%^{12,13}, 1.1 to 44.4%^{14,15} and 8.7 to 44.5%^{16,17}, respectively.

Despite the well-documented strength increases with resistance training in youths, eccentric muscle actions have received little attention in the literature. Although speculative, this might be owing to coach perceptions of eccentric contractions. That is, the high force and lengthening nature of eccentric actions might cause damage to the muscle. However, it has consistently been reported that bouts of eccentric exercise in youths results in similar, and often attenuated, symptoms of exercise induced muscle damage, as compared to adults¹⁸⁻²⁰. When used longitudinally, eccentrically-biased training can provide a range of benefits including enhanced concentric and eccentric strength²¹, change of direction ability^{22,23}, sprint and jump performance^{23,24} and decreased injury incidence²⁴. Moreover, a recent meta-analysis concluded that the Nordic hamstring exercise (NHE) can induce positive changes in muscle architecture and strength²⁵. Notwithstanding the performance related benefits, the NHE has the ability to reduce hamstring injuries by up to 51% in a sample of 8459 athletes²⁶ and is advocated within youth soccer injury prevention programmes²⁷. This is particularly important when considering that hamstring strains account for 12% of injuries among 17 top flight European soccer teams²⁸ and is considered the most important injury risk factor²⁹. Though these injury rates are lower in youths compared to adults, injury prevention is still an important issue in youths as its occurrence negatively affects participation in sport and can have medical implications³⁰. Before the efficacy of such a training study can be investigated, the reliability of the test must first be determined. Consequently, accurate assessment of eccentric hamstring strength via a field-based measure such as the NHE is warranted. Whilst the reliability of the NHE has been determined in adults³¹, there is no available data in youths. Previous work has established that isokinetic eccentric³² and isometric hamstring³³ muscle actions can be reproduced within acceptable limits in males youth (~13 and 17 years, respectively). However, these authors did not ascertain if maturation altered the reliability of these exercises. This is a concern in testing the youth athlete as disrupted motor coordination can occur around the interval of maximal growth³⁴. Thus, it is plausible that the reliability of a test could change as it is used across the maturational spectrum³⁵. A study that determines the reliability of measures of eccentric hamstring strength across maturation stages would help applied practitioners monitor strength adaptations with confidence. Whether for athlete support or inclusion in research, the importance of exercise test reliability is well established^{36,37}. Moreover, by establishing the reliability of an exercise test, a practitioner/research can identify if a change has occurred due to a training intervention rather than biological variation or maturation³⁷. Consequently, the aim of this study is to establish the reliability of the NHE in male youths using a field-based device (NordBord). A further aim was to establish if maturation stage, determined by both chronological age and maturity offset, influences the reliability of the NHE.

1 **Methods**

2 *Subjects*

3 Sixty-four male youth football players aged between 10 and 16 years took part in the study (Table 1).
4 All participants were free from lower-limb musculoskeletal injuries, physically active and participated
5 regularly in association football training. None of the participants were involved in any formalised
6 strength and conditioning programmes and had no prior experience of performing the NHE. Parental
7 informed consent was obtained for the study which was approved by the host institutions ethics
8 committee.

9 [Insert Table 1 about here]

10 11 *Study design*

12 This study employed a repeated measures design in which participants performed the NHE on six
13 separate occasions. Before each performance participants complete a standardised warm up
14 consisting low-intensity jogging, change of direction, jumping tasks and dynamic lower-limb
15 stretching. In the first four sessions, participants were familiarised to the NHE. Participants were
16 deemed 'familiarised' when they could perform multiple repetitions with the correct technique (see
17 below). These familiarisation trials were not used for analysis. For the testing trials participants
18 attended on two occasions, separated by seven days. During each testing trial, participants completed
19 three repetitions of the NHE. Participants did not report any symptoms of exercise-induced muscle
20 damage (e.g. reduced muscle function or elevated muscle soreness).

21 22 *Methodology*

23 *Anthropometry*

24 Age, stature and body mass were obtained prior to testing. Participants' standing and seated height
25 were measured using a stadiometer (Seca Model 213, Birmingham, England). Body mass was
26 measured using a calibrated electronic scale (Seca Model 813, Birmingham, England). Maturity offset
27 was calculated using age, body mass, standing and seated height³⁸. This method provides a practical,
28 non-invasive and accurate measure of maturation status³⁸. Pre-PHV and mid-post PHV participants
29 were categorized as exceeding -2 years and between -1 to +2.5 years, respectively, from PHV. In
30 addition, participants were categorised chronologically by age (i.e. under 11, 13 and 16 years).

31 32 *Eccentric hamstring strength*

33 Eccentric hamstring strength was determined using the NHE on the NordBord (Nordbord, Vald
34 Performance, Australia). The NHE is deemed a reliable marker of peak eccentric hamstring force in
35 adult males athletes (coefficient of variation (CV) % = 5.8 to 8.5%)³¹. Participants were instructed to
36 kneel on the padded part of the NordBord and were positioned with their ankles secured with padded
37 hooks, which were attached to load cells. Participants were positioned so that their ankles were
38 perpendicular to the lower limb and the hooks superior to the lateral malleolus. Participants were
39 instructed to gradually lower their upper-body whilst trying to resistance the movement by
40 contracting the hamstrings. With trunk and hips in a neutral position, participants were encouraged
41 to maintain an upright posture. Coaching cues (i.e. "stay as tall as you can", "slowly fall like a tree")
42 were provided. During the movement, participants arms were flexed at the elbow so that their palms
43 were pronated at shoulder level. In the final stages of the movement participants were allowed to use
44 their hands to buffer their fall. The researchers assisted in returning the participants back to the
45 starting position. Participants performed three repetitions with a self-selected rest that ranged from
46 10 to 15s. The NordBord provides peak forces for each limb thus bilateral peak force was determined
47 by averaging the three scores from each limb. Bilateral peak force was divided by body mass to
48 established relative peak force. The imbalance in peak force between limbs was calculated as the
49 absolute difference between left and right limbs.

50 51 *Statistical analysis*

1 The average value of peak force (N) across the three repetitions was used for analysis. Data was found
2 to be normally distributed according to the Shapiro-Wilk statistic ($P > 0.05$). A paired samples t-test
3 was used to determine the differences in peak force metrics between trials 1 and 2. The reliability of
4 the NHE was quantified via the typical error (TE; standard deviation of the differences divided by $\sqrt{2}$)
5 and CV (TE divided by the grand mean test-retest score, multiplied by 100) statistics³⁹. The smallest
6 worthwhile change (SWC; 0.2 multiplied by the shared standard deviation) was calculated to provide
7 a practical interpretation of the findings. The dependent variables were considered capable of
8 detecting small changes if the TE was less than the SWC⁴⁰.

9 10 **Results**

11 The descriptive characteristics for the NHE across groups and for the entire sample is presented in
12 table 2. A paired samples t-test revealed a significant bias from trial 1 to trial 2 for 25 of the 30
13 comparisons ($P < 0.05$). For all dependent variables and for all comparisons, the TE was greater than
14 the SWC (Table 3). Across the age groups, the CVs for left (CV% = 5.6 to 7.4), right (CV% = 5.9 to 8.7),
15 bilateral (CV% = 6.1 to 7.4) relative peak force (CV% = 6.3 to 6.6) were generally favourable and
16 revealed no clear trend in agreement. The peak force imbalance between left and right limbs
17 demonstrate poor agreement between trials (CV% = 33.1 to 38.3) across the age groups. Reliability
18 for those pre-PHV was better than those mid-post-PHV for left (CV% = 5.7 vs 6.9, respectively), right
19 (CV% = 4.8 vs 8.5, respectively), bilateral (CV% = 4.9 vs 7.3, respectively), relative peak force (CV% =
20 5.0 vs 7.2, respectively) and the imbalance (CV% = 35.3 vs 36.1, respectively).

21
22 [Insert Tables 2 and 3 about here]

23 24 **Discussion**

25 We sought to establish the reliability of the Nordic hamstring exercise using a field-based device in
26 male youth soccer players. We report that certain measures of peak force (i.e. individual limits,
27 bilateral and relative force) can be reproduced within acceptable limits for this population. A
28 secondary aim was to investigate if maturation affected the reliability of the NHE. Whilst reliability
29 was not different across chronological age groups, those classified as pre-PHV demonstrated better
30 agreement between trials than those in the mid-post-PHV group.

31 Atkinson and Nevill³⁶ propose that the reliability of a measure/test is dependent on the
32 setting that it is applied in. For youths who resistance train, lower-body strength (maximal force,
33 torque or kilograms) can increase by up to 44.5%¹²⁻¹⁷ depending on maturation stage. As such a
34 variation of 10% (i.e. a CV of 10%) would allow such improvements to be detected. Importantly, the
35 TE and SWC can be incorporated to facilitate the analysis³⁷.

36 The reliability for the whole sample, for left, right, bilateral and relative peak force was
37 generally favourable (CV% = 6.3 to 8.3), albeit none of the TEs were able to detect the SWC. Previous
38 work has also reported good reliability for eccentric hamstring exercise in circumpubertal males ($r =$
39 0.71 to 0.85)³² and isometric hamstring dynamometry in male youths (minimal detectable change =
40 11.8 to 15.9%)³³. Similarly, during the NHE (in adult males) Opar and colleagues³¹ reported low CVs
41 for peak force (5.8 to 8.5%). However, our study adds to the current body of literature by
42 demonstrating poor reliability of lower-limb strength asymmetry during the NHE (TE and CV of 9.1N
43 and 36.9%, respectively). It is unclear why such poor reliability was observed, especially given the good
44 agreement observed for left and right limbs individually. Irrespective of the mechanism, this data
45 indicates that when assessing muscular imbalances during the NHE across a range of ages, applied
46 practitioners should be cautious.

47 When categorising the participants by chronological age, reliability was similar across U11
48 (CV% = 5.9 to 7.4%), U13 (CV% = 5.6 to 7.8) and U16 (CV% = 6.6 to 8.7) groups, although none of the
49 TE were lower than the SWC. Conversely, when categorised by maturity offset those in pre-PHV group
50 demonstrated better reliability (TE = 0.22 to 9.3N, CV% = 4.8 to 35.5) than the mid-post-PHV group
51 (TE = 0.30 to 22.5, CV% = 6.9 to 36.1). The reliability observed for these maturity groups can

1 comfortably detect the increases in strength that occur in those pre- (~36%)¹³ and mid-post-PHV
2 (~44.5%)¹⁵. However, practitioners should adopt caution when establishing muscular imbalances
3 using the NHE across maturation stages given the large random errors (TE = > 8.1 N and CVs > 35.5%).
4 That we observed better reliability in those pre-PHV than their more mature counterparts, might be
5 due to the well-established disruptions in motor performance that occur during maturation³⁴.
6 Moreover, these data reinforce the importance of categorising youths by maturation rather than
7 chronological age^{11,41}. Nonetheless, the reliability of left and right limb, bilateral and relative peak
8 force for the mid-post-PHV group was still acceptable and thus typical changes in strength can still be
9 detected.

10 In the present study, we observed a systematic bias for several of the dependent variables.
11 While the reasons for this are not entirely clear, the larger values in trial 2, than trial 1, might be
12 indicative of short-term adaptation to the exercise. That participants were given four familiarisation
13 attempts before the testing trials and could competently perform the exercise might support this.
14 Nonetheless, applied practitioners should consider these short-term changes when assessing peak
15 force variables using the NHE.

16 **Conclusion**

17 To our knowledge, this is the first study to provide a comprehensive assessment of Nordic hamstring
18 exercise reliability in male youth soccer players. We report that, despite not being able to detect the
19 small changes, the reliability of the exercise is generally favourable. Notably, the reliability in less
20 mature (i.e. pre-PHV) participants was generally better than their mid-post-PHV counterparts.
21 Nonetheless, applied practitioners can be confident in assessing changes in eccentric hamstring
22 strength using the NHE. However, when assessing muscular imbalances using the NHE, applied
23 practitioners should exercise caution given the large random errors.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **REFERENCES**

- 1 1. Knuttegen HG, Komi P V. Strength and Power in Sport. In: Komi P V., ed. *Strength and Power in Sport*. 2nd ed. Oxford, UK: Blackwell Publishing Company; 2003:3-11.
- 2 2. Duchateau J, Enoka RM. Neural control of lengthening contractions. *J Exp Biol*. 2016;219(2):197-204.
- 3 3. Lloyd RS, Faigenbaum AD, Stone MH, et al. Position statement on youth resistance training: The 2014 International Consensus. *Br J Sports Med*. 2014;48(7):498-505.
- 4 4. Comfort P, Stewart A, Bloom L, Clarkson B. Relationship between strength sprint and jump performance in well-trained youth soccer players. *J Strength Cond Res*. 2014;28(1):173-177.
- 5 5. Branta C, Haubenstricker J, Seefeldt V. Age changes in motor skills during childhood and adolescence. *Exerc Sport Sci Rev*. 1984;12(1):467-520.
- 6 6. Kraemer WJ, Fry AC, Conroy B, Frykman PN, Hoffman J. Resistance training and youth. *Pediatr Exerc Sci*. 1989;1(4):336-350.
- 7 7. Granacher U, Goesele A, Roggo K, et al. Effects and mechanisms of strength training in children. *Int J Sports Med*. 2011;32:357-364.
- 8 8. Ramsay JA, Blimkie CJR, Smith K, Garner S, MacDougall JD, Sale DG. Strength training effects in prepubescent boys. *Med Sci Sports Exerc*. 1990;22(5):605-614.
- 9 9. Tonson A, Ratel S, Fur Y Le, Cozzone P, Bendahan D. Effect of maturation on the relationship between muscle size and force production. *Med Sci Sports Exerc*. 2008;40(5):918-925.
- 10 10. Moran J, Sandercock GRH, Ramírez-Campillo R, Meylan C, Collison J, Parry DA. A meta-analysis of maturation-related variation in adolescent boy athletes' adaptations to short-term resistance training. *J Sports Sci*. 2016;35(11):1041-1051.
- 11 11. Moran J, Sandercock GRH, Ramirez-Campillo R, et al. Maturation-related differences in adaptations to resistance training in young male swimmers. *J Strength Cond Res*. 2018;32(1):139-149.
- 12 12. Meylan CMP, Cronin JB, Oliver JL, Hopkins WG, Contreras B. The effect of maturation on adaptations to strength training and detraining in 11-15-year-olds. *Scand J Med Sci Sport*. 2014;24(3):156-164.
- 13 13. Chaouachi A, Othman AB, Hammami R, Drinkwater EJ, Behm DG. The combination of plyometric and balance training improves sprint and shuttle run performances more often than plyometric-only training with children. *J Strength Cond Res*. 2014;28(2):401-412.
- 14 14. Channell BTC, Barfield JP. Effect of Olympic and traditional resistance training on vertical jump improvement in high school boys. *J Strength Cond Res*. 2008;22(11):1522-1527.
- 15 15. Szymanski DJ, Szymanski JM, Molloy JM, Pascoe DD. Effect of 12 weeks of wrist and forearm training on high school baseball players. *J Strength Cond Res*. 2004;18(3):432-440.
- 16 16. Kotzamanidis C, Chatzopoulos D, Michailidis C, Papaikovou G, Patikas D. The effect of a combined high-intensity strength and speed training program on the running and jumping ability of soccer players. *J Strength Cond Res*. 2005;19(2):369-375.
- 17 17. Harries SK, Lubans DR, Callister R. Comparison of resistance training progression models on maximal strength in sub-elite adolescent rugby union players. *J Sci Med Sport*. 2016;19(2):163-169.
- 18 18. Gorianovas G, Skurvydas A, Streckis V, Brazaitis M, Kamandulis S, McHugh MP. Repeated bout effect was more expressed in young adult males than in elderly males and boys. *Biomed Res Int*. 2013.
- 19 19. Pullinen T, Mero A, Huttunen P, Pakarinen A, Komi P V. Resistance exercise-induced hormonal response under the influence of delayed onset muscle soreness in men and boys. *Scand J Med Sci Sport*. 2011;21:184-194.
- 20 20. Deli CK, Fatouros IG, Paschalis V, Avloniti A. A comparison of exercise-induced muscle damage following maximal eccentric contractions in men and boys. *Pediatr Exerc Sci*. 2017;29:316-326.
- 21 21. Walker S, Blazeovich AJ, Haff GG, Tufano JJ, Newton RU, Häkkinen K. Greater strength gains after training with accentuated eccentric than traditional isoinertial loads in already strength-trained men. *Front Physiol*. 2016;7:1-12.

- 1 22. de Hoyo M, Sañudo B, Carrasco L, et al. Effects of 10-week eccentric overload training on kinetic
2 parameters during change of direction in football players. *J Sports Sci.* 2016;34(14):1380-1387.
- 3 23. de Hoyo M, de la Torre A, Pradas F, et al. Effects of eccentric overload bout on change of
4 direction and performance in soccer players. *Int J Sports Med.* 2015;36:308-314.
- 5 24. de Hoyo M, Pozzo M, Sañudo B, et al. Effects of a 10-week in-season eccentric-overload training
6 program on muscle-injury prevention and performance in junior elite soccer players. *Int J*
7 *Sports Physiol Perform.* 2015;10:46-52.
- 8 25. Cuthbert M, Ripley N, McMahon JJ, Evans M, Haff GG, Comfort P. The effect of Nordic
9 hamstring exercise intervention volume on eccentric strength and muscle architecture
10 adaptations: A systematic review and meta-analyses. *Sport Med.* 2019;EPUB.
11 doi:10.1007/s40279-019-01178-7
- 12 26. Van Dyk N, Behan FP, Whiteley R. Including the Nordic hamstring exercise in injury prevention
13 programmes halves the rate of hamstring injuries: A systematic review and meta-analysis of
14 8459 athletes. *Br J Sports Med.* 2019;(August 2018):1-10.
- 15 27. Owoeye OBA, Akinbo SRA, Tella BA, Olawale OA. Efficacy of the FIFA 11+ Warm-Up Programme
16 in Male Youth Football: A Cluste...: EBSCOhost. *J Sport Sci Med.* 2014;(December 2013):321-
17 328.
18 [http://web.a.ebscohost.com.ezproxy.library.unlv.edu/ehost/pdfviewer/pdfviewer?vid=3&sid](http://web.a.ebscohost.com.ezproxy.library.unlv.edu/ehost/pdfviewer/pdfviewer?vid=3&sid=66859ceb-8540-4bf1-a8c3-dc4fb95c5ac6%40sessionmgr4002&hid=4106)
19 [=66859ceb-8540-4bf1-a8c3-dc4fb95c5ac6%40sessionmgr4002&hid=4106.](http://web.a.ebscohost.com.ezproxy.library.unlv.edu/ehost/pdfviewer/pdfviewer?vid=3&sid=66859ceb-8540-4bf1-a8c3-dc4fb95c5ac6%40sessionmgr4002&hid=4106)
- 20 28. Ekstrand J, Hägglund M, Waldén M. Injury incidence and injury patterns in professional
21 football: The UEFA injury study. *Br J Sports Med.* 2011;45(7):553-558.
- 22 29. Read PJ, Jimenez P, Oliver JL, Lloyd RS. Injury prevention in male youth soccer: Current
23 practices and perceptions of practitioners working at elite English academies. *J Sports Sci.*
24 2018;36(12):1423-1431. [https://doi.org/10.1080/02640414.2017.1389515.](https://doi.org/10.1080/02640414.2017.1389515)
- 25 30. Valle X, Malliaropoulos N, Párraga Botero JD, et al. Hamstring and other thigh injuries in
26 children and young athletes. *Scand J Med Sci Sport.* 2018;28(12):2630-2637.
- 27 31. Opar DA, Piatrkowski T, Williams MD, Shield A. A novel device using the Nordic hamstring
28 exercise to assess eccentric knee flexor strength: A reliability and retrospective injury study. *J*
29 *Orthop Sports Phys Ther.* 2013;43(9):636-640.
- 30 32. Kellis E, Kellis S, Gerodimos V, Manou V. Reliability of isokinetic concentric and eccentric
31 strength in circumpubertal soccer players. *Pediatr Exerc Sci.* 1999;11:218-228.
- 32 33. Wollin M, Purdam C, Drew MK. Reliability of externally fixed dynamometry hamstring strength
33 testing in elite youth football players. *J Sci Med Sport.* 2016;19(1):93-96.
- 34 34. Quatman-Yates CC, Quatman CE, Meszaros AJ, Paterno M V, Hewett TE. A systematic review
35 of sensorimotor function during adolescence: A developmental stage of increased motor
36 awkwardness? *Br J Sports Med.* 2012;46(9):649-655.
- 37 35. Moeskops S, Oliver JL, Read PJ, et al. Within- and Between-Session Reliability of the Isometric
38 Midthigh Pull in Young Female Athletes. *J Strength Cond Res.* 2018;32(7):1892-1901.
- 39 36. Atkinson G, Nevill A. Statistical methods for assssing measurement error (reliability) in
40 variables relevant to sports medicine. *Sport Med.* 1998;26(4):217-238.
- 41 37. Fernandes JFT, Lamb KL, Twist C. The intra- and inter-day reproducibility of the FitroDyne as a
42 measure of multi-jointed muscle function. *Isokinet Exerc Sci.* 2016;24(1):39-49.
- 43 38. Mirwald RL, Baxter-Jones ADG, Bailey DA, Beunen GP. An assessment of maturity from
44 anthropometric measurements. *Med Sci Sports Exerc.* 2002;34(4):689-694.
- 45 39. Hopkins WG. Measures of reliability in sports medicine and science. *Sport Med.*
46 2000;30(5):375-381.
- 47 40. Pyne DB. Interpreting the results of fitness testing. In: *International Science and Football*
48 *Symposium.* ; 2003:1-6.
- 49 41. Moran J, Sandercock G, Clark CCT, Fernandes JFT, Drury B. A meta-analysis of resistance
50 training in female youth: Its effect on muscular strength, and shortcomings in the literature.
51 *Sport Med.* 2018.
- 52

1 **Table 1.** Anthropometric characteristics of the participants
2

	All (n=64)	U11 (n=17)	U13 (n=29)	U16 (n=18)	Pre-PHV (n=29)	Mid-post-PHV (n=35)
<i>Age (y)</i>	13.2 ± 1.7	10.8 ± 0.3	12.1 ± 0.7	15.4 ± 0.4	11.6 ± 0.9	14.5 ± 0.9
<i>Mass (kg)</i>	52.3 ± 12.8	42.0 ± 3.7	39.6 ± 2.5	63.7 ± 6.6	40.0 ± 3.8	62.3 ± 7.2
<i>Stature (cm)</i>	161.9 ± 13.4	148.6 ± 2.8	149.2 ± 4.7	173.9 ± 6.7	148.4 ± 5.1	172.4 ± 6.7

3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

1 **Table 2.** Mean \pm standard deviations values for left, right and bilateral peak flexor force, relative peak force and peak force imbalances during the Nordic
 2 hamstring exercise
 3

	Left limb (N)		Right limb (N)		Bilateral force (N)		Relative force (N \cdot kg)		Imbalance (N)	
	Trial 1	Trial 2	Trial 1	Trial 2	Trial 1	Trial 2	Trial 1	Trial 2	Trial 1	Trial 2
U11	141.7 \pm 42.8	159.1 \pm 44.7*	165.2 \pm 41.8	176.1 \pm 40.2*	153.5 \pm 40.3	167.6 \pm 41.7*	3.9 \pm 0.9	4.2 \pm 1.0*	29.5 \pm 17.7	20.5 \pm 10.3
U13	211.8 \pm 51.7	221.3 \pm 55.7	220.0 \pm 52.1	240.4 \pm 63.2*	215.9 \pm 51.1	230.9 \pm 58.6*	4.3 \pm 0.7	4.5 \pm 0.7*	18.8 \pm 15.6	24.3 \pm 14.7*
U16	244.5 \pm 65.3	264.7 \pm 54.6*	260.4 \pm 55.5	296.1 \pm 59.8*	252.5 \pm 58.9	280.4 \pm 56.0*	4.0 \pm 1.1	4.4 \pm 1.1*	25.5 \pm 18.9	32.4 \pm 22.7
Pre-PHV	157.8 \pm 41.8	168.4 \pm 37.5*	175.7 \pm 39.5	183.5 \pm 36.4*	166.7 \pm 38.7	176.0 \pm 35.9*	4.2 \pm 0.9	4.4 \pm 0.9*	25.4 \pm 17.0	20.3 \pm 11.0
Mid-post-PHV	234.9 \pm 64.8	252.4 \pm 61.5*	247.5 \pm 57.8	280.2 \pm 66.0*	241.2 \pm 60.0	266.3 \pm 62.8*	4.0 \pm 0.9	4.4 \pm 0.9*	22.8 \pm 18.3	29.9 \pm 19.3*
All	202.4 \pm 66.2	217.0 \pm 65.0*	216.8 \pm 61.2	239.0 \pm 71.8*	209.6 \pm 62.5	228.0 \pm 67.7*	4.1 \pm 0.9	4.4 \pm 0.9*	23.5 \pm 17.5	25.6 \pm 16.8

4 *denotes significantly different between trials 1 and 2 ($P < 0.05$).
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24

1 **Table 3.** Reliability statistics for left, right and bilateral peak flexor force, relative peak force and strength imbalances during the Nordic hamstring exercise

2

		Left limb	Right limb	Bilateral force	Relative force	Imbalance
All	<i>TE (N)</i>	14.0	18.9	15.4	0.27	9.1
	<i>SWC (N)</i>	4.0	5.3	4.4	0.08	2.6
	<i>CV (%)</i>	6.7	8.3	7.0	6.3	36.9
U11	<i>TE (N)</i>	11.1	10.0	9.8	0.26	9.6
	<i>SWC (N)</i>	3.1	2.8	2.8	0.07	2.7
	<i>CV (%)</i>	7.4	5.9	6.1	6.3	38.3
U13	<i>TE (N)</i>	12.1	17.9	14.1	0.28	7.1
	<i>SWC (N)</i>	3.4	5.1	4.0	0.08	2.0
	<i>CV (%)</i>	5.6	7.8	6.3	6.3	33.1
U16	<i>TE (N)</i>	18.3	24.3	19.8	0.28	10.8
	<i>SWC (N)</i>	5.2	6.9	5.6	0.08	3.1
	<i>CV (%)</i>	7.2	8.7	7.4	6.6	37.3
Pre-PHV	<i>TE (N)</i>	9.3	8.7	8.4	0.22	8.1
	<i>SWC (N)</i>	2.6	2.5	2.4	0.06	2.3
	<i>CV (%)</i>	5.7	4.8	4.9	5.0	35.5
Mid-post-PHV	<i>TE (N)</i>	16.7	22.5	18.4	0.30	9.5
	<i>SWC (N)</i>	4.7	6.4	5.2	0.09	2.7
	<i>CV (%)</i>	6.9	8.5	7.3	7.2	36.1

3

