

**Knowledge of results during vertical jump testing: an effective method to increase the performance but not the consistency of vertical jumps**

Garcia-Ramos, Amador; Janicijevic, Danica; Cobo-Font, Juande; Marcos-Frutos, Daniel; Fernandes, John; Taube, Wolfgang; Pérez-Castilla, Alejandro

*Published in:*  
Sports Biomechanics

*Publication date:*  
2020

*The re-use license for this item is:*  
CC BY-NC-ND

*This document version is the:*  
Peer reviewed version

*The final published version is available direct from the publisher website at:*  
[10.1080/14763141.2020.1764090](https://doi.org/10.1080/14763141.2020.1764090)

**[Find this output at Hartpury Pure](#)**

*Citation for published version (APA):*  
Garcia-Ramos, A., Janicijevic, D., Cobo-Font, J., Marcos-Frutos, D., Fernandes, J., Taube, W., & Pérez-Castilla, A. (2020). Knowledge of results during vertical jump testing: an effective method to increase the performance but not the consistency of vertical jumps. *Sports Biomechanics*. <https://doi.org/10.1080/14763141.2020.1764090>

# Title: Individual vs. Group-based Strategies for Weight Loss and Management in Adults: Pen Profile Perceptions

Cain C. T. Clark<sup>1\*</sup>, Amie Chambers<sup>2</sup>, John F. T. Fernandes<sup>2</sup>, Martin Musalek<sup>3</sup>, Robert Podstawski<sup>4</sup>, Emma L. J Eyre<sup>1</sup>, Michael Duncan<sup>1</sup>

<sup>1</sup> Centre for Sport, Exercise and Life Sciences, Coventry University, Coventry (U.K.), CV1 5FB

<sup>2</sup> Hartpury University, Applied Sport Science, Gloucestershire (U.K.), GL19 3BE.

<sup>3</sup> Faculty of Physical Education and Sport, Charles University, Prague, Czech Republic.

<sup>4</sup> University of Warmia and Mazury, Olsztyn, Department of Physical Education and Sport Olsztyn, Poland

\*Corresponding author: Dr. Cain C. T. Clark., [Cain.Clark@coventry.ac.uk](mailto:Cain.Clark@coventry.ac.uk); Twitter: @DrCainCTClark;

ORCID: <http://orcid.org/0000-0002-6610-4617>

## Abstract

### *Aims*

The aim of this study was to explore perceptions of barriers, facilitators, strategies and successes in individual vs. group-based weight management programmes.

### *Methods*

Forty-two, overweight (Body Mass Index  $\geq 25$ -kg·m<sup>2</sup>) participants (aged 32-63y) volunteered to take part in this study. All participants completed a 3-month weight loss programme, randomised to a group-based (n=21), or self-monitoring only (n=21) approach, respectively. Participants subsequently participated in a semi-structured interview (60±7 mins) to explore individual and collective perceptions of barriers, facilitators, strategies and successes.

### *Results*

Convergent themes were found for individual and group strategies for facilitators, strategies and successes. Divergent themes were found between groups for barriers, group participants highlighted expense of commercial products, and knowledge of nutrition and dieting, whilst individual participants reported (lack of) social support from peers, (lack of) motivation, and occupation.

### *Conclusion*

Key stakeholders, facilitators, and individuals must consider these factors prior to the advocacy any weight loss strategy.

Key words: Obesity; Overweight; Weight-loss; Management; Strategy

## 32 **Introduction**

33 Obesity, broadly speaking, is characterised by having a body-mass index above 30 kg·m<sup>2</sup>, and  
34 described as a non-communicable risk factor (Atay & Bereket, 2016). The proportion of adults in the United  
35 Kingdom (UK) that are overweight or obese has risen from 57.6% to 68% in men, and from 48.6% to 58%  
36 in women between 1993 and 2018; representing an estimated total economic burden of £27 billion (Clark,  
37 2018). Indeed, obesity in the UK (and worldwide) is acknowledged as to be an epidemic (Clark, 2018);  
38 whilst physical activity has been identified as an integral contributor to a healthy lifestyle (Saunders et al,  
39 2016) and can provide immediate and future health benefits (Telema et al, 2013; Shiri et al, 2013), physical  
40 inactivity is the largest contributor to risk factors for non-communicable diseases worldwide (WHO, 2010),  
41 exacerbating the prevalence of obesity. Strong relationships exist between physical activity and health, with  
42 higher physical activity levels leading to reduced risks of coronary heart disease (Li et al, 2012),  
43 hypertension (Peters et al, 2006), non-insulin dependent diabetes mellitus (LaMonte, Blair & Church, 2005),  
44 stroke (Goldstein, 2010), colon cancer (Wolin et al, 2009), osteoporotic fractures (de Kam et al, 2009) and  
45 depression (Martinsen, 2008). Notwithstanding, weight loss and maintenance is multi-factorial, and it has  
46 become increasingly apparent that nutrition, support and encouragement, among other facilitators, are  
47 pivotal tenets of any weight loss strategy.

48  
49 The benefits of regular physical activity and a healthy lifestyle have been clearly set out across the  
50 life course (WHO, 2010). As such, it has become common practice to promote and encourage overweight  
51 and obese individuals towards weight loss strategies and groups, yet there remains no consensus on the most  
52 efficacious weight loss strategy. Weight loss or maintenance strategies can be broadly dichotomised into  
53 self-monitoring or group-based. Self-monitoring is a common, very easy method of weight loss. Originally,  
54 this method was observed using paper records of an individual's diet (Sperduto, Thompson and O'Brien,  
55 1986). Such self-monitoring techniques have further expanded to include the addition of physical activity  
56 and how individuals pair both factors as a method of behaviour change to lose weight (Wadden, Butryn and  
57 Wilson, 2007). Additionally, self-monitoring in terms of self-weighing to track progress has also been

58 identified (Linde et al. 2005). The self-reliant nature of this strategy is asserted to make the individual more  
59 aware of their current behaviours; making goal setting easier, enabling them to track progress and then put  
60 a number to this progress with the weighing (Foster, Makris and Bailer, 2005).

61  
62 Group-based interventions comprise an alternative to self-monitoring that is equally popular as a  
63 method of weight loss. For example, the commercially prevalent Weight Watchers (WW) or Slimming  
64 World (SW), which are often referrals from primary health care practitioners. These are commercial weight-  
65 loss programmes that individuals can sign up to of their own volition or are often referrals from primary  
66 health care practitioners (Jolly et al., 2007). Individuals sign up with this programme and attend sociable  
67 meetings to have a weigh in once a week; and are also able to purchase food, drinks and snacks sold by  
68 these companies all in aid of losing weight.

69  
70 Empirical evidence has shown the structure and intensity of contact within a weight loss method  
71 are significantly correlated with success (Jensen *et al.*, 2013; Wadden *et al.*, 2011); however, inter-  
72 individual interpretation of such key terms is equivocal. Metzgar *et al.* (2014) reported that accountability  
73 and support, exercise, motivation, total lifestyle change and eating patterns majorly influenced weight loss  
74 maintenance. Additionally, Elfhag and Rossner (2005) identified factors associated with successful weight  
75 loss included; a high amount of initial weight loss, goal setting, physical activity, regular eating patterns and  
76 controlled eating habits. Furthermore, to enhance weight maintenance specifically Elfhag and Rossner  
77 (2005) asserted that individuals look for an internal motivation to lose weight, high self-efficacy, coping  
78 strategies, social support and strong psychological strength. To ensure this, they suggested that a weight  
79 loss method requires an appropriate amount of guidance and support to ensure weight maintenance, which  
80 is of high importance within the first three months.

81  
82 Recently, Lemstra and colleagues (2016) observed that self-monitoring weight loss programs have  
83 lower adherence than group-based programmes, 41.5% vs. 68.6%, respectively. There are innumerable

84 tenets of a successful and sustained weight loss and management programme, yet inter-individual  
85 preferences and perceptions of weight-management strategies are less well known; where if such  
86 perceptions of weight-management strategies might be best elucidated via qualitative research methodology  
87 as it provides more in depth understanding of inter-individual factors that may predispose a weight  
88 management strategy to be more or less effective. Therefore, the aim of this study was to explore perceptions  
89 of barriers, facilitators, strategies and successes in individual versus group-based weight loss programmes.

90

## 91 **Methods**

### 92 *Overview*

93 This study drew on data collected via interviews conducted with forty-two overweight (Body Mass  
94 Index  $\geq 25$  kg.m<sup>2</sup>) participants (aged 32-63 years) at the conclusion of a 3-month weight loss programme,  
95 randomised to a commercially prevalent group-based (n=21), or self-monitoring only (n=21) approach,  
96 respectively. The project received institutional ethical approval and conformed to the Declaration of  
97 Helsinki.

98

### 99 *Participants and Settings*

100 Forty-two overweight (Body Mass Index  $\geq 25$ -kg.m<sup>2</sup>) participants (aged 32-63y) volunteered to take  
101 part in this study. Following anamnesis questioning; all participants verbally confirmed no prior experience  
102 of formally taking part in a weight-loss plan, group, or strategy, so to avoid prior personal experiences  
103 influencing opinions and beliefs, *a priori*. Participants were randomized into either 3-months of self-  
104 monitoring weight-loss only, or a commercial weight-loss group.

105

106 Demographic information (age and sex) were collected via Web-based survey, completed during  
107 the first week of the study. At the conclusion of the 3-month period, all individuals participated in a semi-  
108 structured interview. A qualitative approach was used to respect the expert knowledge of the participants  
109 and to enable them to provide insights into their experiences (Ridgers et al, 2012). The interviews followed

110 a semi-structured format and were designed to address individual perceptions of barriers, facilitators,  
111 strategies and successes; related to respective weight loss strategies. In total, 42 semi-structured interviews  
112 (60±7 mins) were conducted in participants' home environment by two of the authors, and digitally  
113 recorded. Interviews were subsequently transcribed verbatim, resulting in 210 pages of raw transcription for  
114 further analyses.

115

### 116 *Data Analyses*

117 Pen profiles were constructed from verbatim transcripts using a manual protocol (see; Mackintosh  
118 et al, 2011; Ridgers et al, 2012). Pen profiles are an increasingly utilized technique that are used to present  
119 analysis outcomes via diagrams of composite key emergent themes, and is considered appropriate and  
120 accessible to researchers with an affinity for both qualitative and quantitative backgrounds. Example,  
121 representative, verbatim quotations were extracted directly from the transcripts to further contextualize the  
122 theme. To provide an indication of the prevalence of the themes, the number of times a specific theme was  
123 mentioned across all interview data is also presented (Ridgers et al, 2012). Consistent with recommended  
124 approaches (Burnard, 1991) one researcher (AC) initially read and analysed the transcripts. These findings  
125 were then presented to another researcher, by means of cooperative triangulation. Having independently  
126 analysed the transcripts, CC then critically questioned the presented thematic analyses and challenged  
127 differing interpretations. A third researcher (RP) subsequently analysed the data in reverse from the pen  
128 profiles back to the transcripts. This process assured the reliability of the data obtained (Ridgers et al, 2012).  
129 Finally, the pen profiles were re-presented to the lead author, who further critically challenged the data. This  
130 process allowed authors to offer alternative interpretations and interrogate the data until a consensus was  
131 reached. Overall, methodological rigor (i.e., credibility and transferability) was demonstrated through  
132 verbatim transcription of data and triangular consensus procedures. Moreover, dependability was  
133 demonstrated through the comparison of pen profiles with verbatim citations and the triangular consensus  
134 processes.

135

136 **Results**

137

138 ***Barriers to weight loss success***

139 Perceptions of barriers to weight loss success were found to be divergent between strategies, where  
140 only time constraints was found as a shared theme. Expense of commercial products, and knowledge of  
141 nutrition and dieting were attributed to the group-based strategy. Less common barriers were weight  
142 maintenance, laziness, the influence of not liking cooking (Figure 1). Exemplar views around barriers  
143 include; “I wasn’t bought up with the understanding of healthy food” (F32), and, “My eating pattern was  
144 very up and down because of my job” (M34). Within the self-monitoring approach, the themes identified  
145 were different to the group intervention participants. However, there were also two common barriers specific  
146 to the weight-loss method; lack of freedom self-monitoring provides, and a lack of social support from  
147 peers. Lesser referred to barriers identified were lack of motivation, the influence of the participants’  
148 occupation (Figure 1).

149

150 **\*\*\*Figure 1 about here\*\*\***

151

152 ***Contributors to weight loss success***

153 Convergent overarching themes were noted for individual and group strategies; including exercise, diet and  
154 personal factors. The group intervention participants identified enjoyment and improving health as  
155 successors to exercise (Figure 2). With respect to diet, reflection, organisation and prior dieting experience  
156 were considered to be important to success (Figure 2). Finally, the personal skills identified were increasing  
157 self-confidence, improving health, being self-motivated, mental state, making the appropriate changes and  
158 self-awareness (Figure 2). The self-monitors observed were improving health, enjoyment, meeting physique  
159 goals and the influence of their occupation to be important to successors to exercise (Figure 3). Convergent  
160 themes were established between the two groups, for diet: organisation, influence of cooking skills,  
161 conscious decision making and reflection; for personal factors: making appropriate changes, self-control,

162 mental state, the influence of their knowledge, time management, influence of their occupation and  
163 improving health; for exercise: health improvement, enjoyment and physique goals were reported (Figure  
164 3). For instance, the self-monitoring group highlighted that their social support comes from friends and  
165 family, “Friends and family are support” (F 41), whilst the group intervention participants stated that their  
166 social support came from the group itself, for example “My support is in the group” (F 44).

167

168

169 \*\*\* **Figure 2 about here** \*\*\*

170

171 \*\*\* **Figure 3 about here** \*\*\*

172

### 173 *Strategy choice influences*

174 Views on strategy choice influences were convergent as overarching themes, where; freedom,  
175 enjoyment, ease, structure and consistent weight loss were evident. For example; “It’s just so easy for me  
176 with the baby” (F34). For the group-based strategy, peer support, weekly schedule and guidance of  
177 improving knowledge were additionally found. Whilst, for the self-monitoring strategy, social support from  
178 friends was additionally referred to (Figure 4).

179

180 \*\*\* **Figure 4 about here** \*\*\*

## 181 **Discussion**

182

183 The aim of this study was to explore perceptions of barriers, facilitators, strategies and successes in  
184 individual only vs. group-based weight management programmes. In accord with our aim, divergent themes  
185 were found between groups for barriers, whilst convergent overarching themes were noted for individual  
186 and group strategies for facilitators, strategies and successes; including exercise, diet and personal factors.  
187 The data presented here offer unique insight into the facets of individual or group-based weight loss



188 strategies which may predispose individuals to be more or less successful when undertaking a weight loss  
189 programme. Such information has yet to be provided by prior work and, as such, the research presented here  
190 extends knowledge in the area/constitutes a novel addition to the area. The following topics will be discussed  
191 as result of the themes found; strategy choice influences, the importance of social support, maintenance of  
192 motivation, organisation, contributors to weight-loss success, guidance, personal factors, exercise, diet,  
193 barriers to weight-loss success, lifestyle and personal factors, and, disadvantages of weight-loss methods.

194

### 195 *Strategy choice influences*

#### 196 *The importance of social support*

197 Social support was deemed to be an important factor for both weight management programmes,  
198 concomitant with previous research, which highlights social support to be a key positive factor contributing  
199 adherence (Lemstra et al. 2016). Whilst comparable, in an overarching view, the present work suggests that  
200 the two groups found social support from different places. Karfopoulou et al. (2016) asserts that although  
201 social support is important, the type of support received can affect weight maintenance. That is, when  
202 comparing individuals who had lost or maintained weight for a year to those who re-gained it, those who  
203 maintained weight loss had received compliments, in comparison to the re-gainers, who had received verbal  
204 instructions. This highlights that the support from peers needs to be positive for the individual to maintain  
205 motivation. This could explain why the self-monitors sought social support within a weight loss method,  
206 given it is not as readily available compared to group intervention.

207

#### 208 *Maintenance of motivation*

209 Other factors relating to maintaining motivation were found through the thematic analyses, where  
210 both the groups acknowledged that a pattern of constant weight loss each week will keep them motivated.  
211 Participants also highlighted their goals in terms of numbers i.e. the importance of monitoring their weight,  
212 to succeed. This could suggest that if the individuals had stopped achieving this weekly weight goal they  
213 may have given up, albeit this does require follow-up work to substantiate. Both self-monitoring and group

214 interventions facilitate participants to track this progress by integrating a weekly weighing/tracking progress  
215 system; although there is a dearth of empirical evidence that specifically identifies this as a mechanism to  
216 motivation maintenance, there is evidence for the underlying principles. For example, Elfhag and Rossner  
217 (2005) support the concept of meeting weight loss goals to be important to participants alongside identifying  
218 that individuals who have a high initial weight loss are more likely to maintain their weight lost. Whilst  
219 Wing and Hill (2001) assert that regular self-monitoring of weight is crucial for weight maintenance,  
220 inferring that if individuals are to choose a weight loss method that incorporates tracking, they are more  
221 likely to maintain their weight; however, how this system is operationalized is somewhat unequivocal.

222  
223         Within the present study, the participants from both groups also conveyed that the weight loss method  
224 they choose needs to provide enjoyment, freedom and ease for them. It is plausible, from the themes  
225 highlighted in this study, that participants seek these from a weight loss method to maintain motivation. The  
226 ease of a weight loss method is crucial, because if it is complicated, or too restrictive and difficult to fit in  
227 to an individuals' lifestyle, it is not conducive to programme maintenance or adherence. Classic empirical  
228 evidence, from Schifter and Ajzen (1985), highlights that idiosyncratic preferences of freedom must be  
229 evident, thereby facilitating enjoyment; without such affordances, motivation will be deleteriously affected.

230

### 231         *Organisation*

232         A further theme identified from both groups constituted the need for a structured eating pattern. This  
233 highlights the importance of self-organisation, which has previously been deemed an important factor to  
234 weight loss success (Elfhag and Rossner, 2005; Kruger, Blanck and Gillespie, 2006). Many of the  
235 participants highlighted how their meal patterns had changed due to the weight loss method they had  
236 adopted. One practical difference between the two groups is that the group intervention, integrated this  
237 structured eating pattern approach from their teachings, for example, using a points-based system (as in  
238 Weight Watchers) to control their intake. Conversely, those that self-monitor are required to organise their  
239 eating patterns for themselves.

240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265

*Guidance*

Within this study, the group intervention participants identified an additional need within their weight loss method, compared to the self-monitors. That is, they highlighted the importance of the organisation of the point-based systems enabling them to control eating habits, in conjunction with the knowledge the leaders or facilitators impart and the provision of educational materials, for example the booklets and recipes. To the authors' knowledge, there exists no research to definitively support that the aforementioned are common influences as to why individuals choose a group-based weight loss intervention. However, within this research, 'guidance' is the difference that dichotomizes the two groups. Notwithstanding, it was evident that self-monitors look for the same core principles within a weight loss method, they are happy volitionally organising their behaviour changes compared to group intervention participants whom seek extra guidance to make appropriate changes.

*Contributors to weight loss success*

*Personal factors*

There are innumerable personality traits and qualities that could influence an individual's likelihood to succeed with weight loss (Montesi et al., 2015), however, within this work, both groups identified similar qualities they deemed important in relation to their weight loss success. Firstly, both groups identified the need to make appropriate changes; whether this be to diet or exercise. Self-control was also identified as an important aspect alongside needing a strong mental state. Finally, both groups identified the importance of improving health, suggesting that if an individual feels intrinsically better, they can see the positive impacts of the weight loss, thereby facilitating increased motivation; a finding which has been affirmed previously (Metzgar et al. 2014). Both groups in the present study also identified other personal factors, albeit in smaller numbers, suggesting the importance of assessing idiosyncratic traits preceding recommendation of a weight loss strategy.

266 *Exercise*

267 The importance of exercise has repeatedly been highlighted for positive health trajectories, whilst  
268 concomitantly being an effective strategy for weight loss; where it is frequently shown to be a predictor of  
269 success in long term healthy weight management (Donnelly et al. 2004; Haus et al. 1994; Ross et al. 2000;  
270 Lejeune et al. 2003; Kahkoska et al. 2018). Empirical evidence is equivocal as to the veracity of diet versus  
271 exercise for weight-loss or maintenance, given they are *two-sides of the same coin*, but increased  
272 effectiveness of using both in combination is globally advocated. In a comprehensive meta-analysis,  
273 Anderson et al. (2001) investigated long-term weight maintenance of participants within structured weight-  
274 loss programmes; six constituent studies concluded that participants who exercised more frequently, had a  
275 significantly greater weight-loss maintenance compared to those who exercised less. Additionally, Wu et  
276 al. (2009) showed that whilst lifestyle interventions with a dietary component result in weight loss,  
277 interventions combining a dietary and physical activity component result in a greater magnitude of weight  
278 loss.

279  
280 Irrespective of weight-loss strategy, those participants engaging in physical activity reported  
281 homogenous themes, highlighting the importance of improving health. Concurrently, participants stated  
282 they enjoyed the forms of exercise they did. Previous research demonstrates adherence rates in exercise to  
283 be increased synchronously with enjoyment (Ryan et al. 1997; Hagberg et al. 2008). There was a range of  
284 physical activities reported, from aqua aerobics, yoga and horse riding to gym-going. Evidently, numerous  
285 forms of exercise appeal to various individuals, suggesting that a symbiotic relationship between weight  
286 loss interventions and proximity to facilities that enable physical activity (e.g. sports and leisure centres)  
287 may be conducive to physical activity engagement; notwithstanding, this is conjecture and necessitates  
288 further investigation. In addition, many self-monitors highlighted that they perceived themselves as having  
289 active jobs, which influenced their decision to exercise or not. Whilst many self-monitors highlight specific  
290 physique goals they wanted to meet in addition to losing weight, such as increasing musculature.

291 *Diet*

292           Diet, concomitant to energy balance, has a major influence on an individuals' weight, thus playing  
293 a key role in weight loss (Bish et al. 2005; Kruger et al. 2004), concomitant to physical activity (referred to  
294 above). Both groups identified the importance of organisation and reflection to their weight loss journey.  
295 There is currently limited empirical data exonerating the concept of self-reflection within weight loss  
296 specifically. However, the concept of self-reflection has been asserted an important skill, which enables an  
297 individual to critically look at what they are doing and make appropriate changes (Baird *et al.* 1991), and  
298 by extension within weight-loss.

299           A stark contrast between the weight loss groups was that the self-monitors identified their cooking  
300 skills as a positive influence. Whilst the group intervention participants highlighted this as a barrier. Kruger  
301 and colleagues (2006) also reported on the importance of organization and meal planning, alongside the  
302 positive influence of liking cooking, as factors relating to successful weight maintenance. This may reveal  
303 a tenet for improvement in group-based strategies, notwithstanding, cooking preference/ability should be  
304 investigated as a potential correlate or determinant of successful weight loss and maintenance.

305

### 306 ***Barriers to weight loss success***

307           Barriers are an acknowledgedly important aspect to consider for potential weight loss; whilst there  
308 are factors individuals cannot control, for example developmental determinants, age and sex, many other,  
309 controllable or changeable, environmental and social factors can heavily influence weight loss and  
310 maintenance, ultimately becoming barriers to success (Weight Management, 2003). Both weight loss groups  
311 identified a range of barriers to weight loss success, however, there were few similarities noted between the  
312 two groups.

313

### 314 ***Lifestyle and personal factors***

315

316           Both groups identified time as a perceived barrier, for example, in response to questions referring to  
317 preparation of food and participating in exercise. In particular, the self-monitors also stated that their

318 occupation was a major time barrier for exercise participation. Much empirical data exists in support of the  
319 current findings (that time was a major barrier), particularly with reference to exercise (Troost et al. 2002;  
320 Andajani-Sutjahjo et al. 2004). Welch et al. (2008) found that 40% of 1,580 women stated time is a barrier  
321 to achieving dietary goals, and 70% of 1,521 women asserted time as a barrier to achieving physical activity  
322 goals.

323  
324 Intrinsic motivation is very common theme across weight loss research, where self-motivation paired  
325 with an internal motivation to lose weight have been identified as predictors of weight loss success (Elfhag  
326 and Rossner, 2005; Teixeira et al. 2005). Within the present study, self-monitors identified (lack of) intrinsic  
327 motivation as a barrier with much greater prevalence than the group intervention, in fact, among group  
328 intervention participants, this was not found as a universal theme. Given the lack of motivation experienced  
329 by only one intervention in the present study, further investigations into this phenomenon should be  
330 encouraged.

331  
332 In the current study, the group intervention participants asserted weight maintenance as an importance  
333 aspect. Highlighting the importance of sustained weight loss or management, and linked to a further  
334 perceived barrier, mental state, which, in a likely cyclical relationship, is exacerbated by reverting towards  
335 starting weight. This relationship warrants deeper investigation to facilitate our understanding of mental  
336 state and weight management.

337  
338 *Disadvantages of weight loss methods*

339  
340 Due to the self-monitoring participants having greater perceived control over their approach/strategy to  
341 weight loss, the barriers they identified were generally not related to the weight loss method itself, rather  
342 the inverse, the abundant freedom was perceived as a potential drawback, whilst paradoxically positively  
343 influencing their choice of strategy. This paradox demonstrates the intricacies of individual weight loss

344 method selection, opposed to a one size fits all approach. Freedom within a weight loss method might  
345 facilitate the participant's control over their physical activity regime, food portions and calorie restrictions.  
346 Whilst these are fundamental tenets of any weight loss programme, it likely necessitates self-control and  
347 understanding (Klem *et al.* 1997; Leahey *et al.* 2014). Conversely, the group intervention participants did  
348 not identify freedom as a barrier, only a strategy choice influence, inferring that the group-based sessions  
349 provide adequate education and support to facilitate better understanding and self-control in their  
350 participants. The group intervention participants highlighted two barriers to success relating to their method  
351 choice. Firstly, the expense of the commercial group products, and, secondly, prior knowledge of nutrition  
352 and dieting. Klohe-Lehman *et al.* (2006) support the importance of knowledge for weight loss success,  
353 highlighting that the amount of weight the participants lost after the intervention, significantly, positively  
354 correlated with the amount of knowledge they gained, albeit subjectively ascertained, this suggests that  
355 greater [perception of] knowledge, is associated with greater weight loss success.

356

### 357 *Limitations*

358

359 This study evaluated participants views in response to 3-months following a weight loss programme,  
360 however, given the importance of weight management, in addition to weight loss, a longer follow up period  
361 could be implemented to assess the changing views, as an individuals' weight loss journey continues. The  
362 interventions employed in this study were 3-months in duration, however, were a longer intervention  
363 implemented, participant views may have differed, and as such, should be considered in further research.  
364 Of importance is that this study did not distinguish between those who were successful or unsuccessful on  
365 their weight loss programme; but assert the information gleaned from this work are equally as important on  
366 ones' weight-loss or maintenance journey, where 3-months is only the beginning. We therefore recommend  
367 that a comprehensive, longitudinal investigation of perceptions and actual weight loss be conducted. Finally,  
368 the participants represented quite a broad age range, it would be advantageous to explore age specific groups  
369 to investigate the intricacies of how views related to weight loss evolve with age.

370

## 371 **Conclusion**

372

373 Whilst facilitators, strategies and successes related to individual only vs group weight-loss approaches  
374 were comparable between groups, divergent perceived barriers were found, highlighting that there is not  
375 one panacea strategy for weight loss or management. Therefore, it is recommended that key stakeholders,  
376 facilitators and individuals must consider these factors prior to the advocacy of any one-particular weight  
377 loss strategy, and use individual/patient experience to facilitate the abatement of perceived barriers.

## 378 **Data Availability**

379 The data used to support the findings of this study are available from the corresponding author upon request

## 380 **Conflicts of Interest**

381 The authors do hereby declare that they have no conflicts of interest relevant to the content of this  
382 manuscript.

383

384

385

386

## **References**

1. Andajani-Sutjahjo, S., Ball, K., Warren, N., Inglis, V. and Crawford, D. (2004) Perceived personal, social and environmental barriers to weight maintenance among young women: a community survey. *International Journal of Behavioral Nutrition and Physical Activity*. 1 (15).
2. Anderson, J.W., Konz, E.C., Frederich, R.C. and Wood, C.L. (2001) Long-term weight-loss maintenance: a meta-analysis of US studies. *American Journal of Clinical Nutrition*. 74 (5), 579-584.



3. Atay, Z., & Bereket, A. (2016). Current status on obesity in childhood and adolescence: Prevalence, etiology, co-morbidities and management. *Obesity Medicine*, 3, 1-9.
4. Baird, J., Fensham, P., Gunstone, R. and White, R. (1991). The importance of reflection in improving science teaching and learning. *Journal of Research in Science Teaching*. 28 (2), 163-182.
5. Bish, C., Blanck, H., Serdula, M., Marcus, M., Kohl, H. and Khan, L. (2005) Diet and Physical Activity Behaviors among Americans Trying to Lose Weight: 2000 Behavioral Risk Factor Surveillance System. *Obesity Research*. 13 (3), 596-607.
6. Burnard P. (1991). A method of analysing interview transcripts in qualitative research. *Nurse Educ Today*, 11(6), 461-466. [Medline: 1775125]
7. Clark, C. C. T. (2017). Is obesity actually non-communicable?. *Ob Med*, 8, 27-28, <https://doi.org/10.1016/j.obmed.2017.10.001>
8. de Kam D, Smulders E, Weerdesteyn V, Smits-Engelsman BC. (2009). Exercise interventions to reduce fall-related fractures and their risk factors in individuals with low bone density: a systematic review of randomized controlled trials. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*, 20(12), 2111-2125.
9. Donnelly, J.E., Smith, B., Jacobsen, D.J., Kirk, E., DuBose, K., Hyder, M., Bailey, B. and Washburn, R. (2004). The role of exercise for weight loss and maintenance. *Best Practice and Research Clinical Gastroenterology*. 18 (6), 1009-1029.
10. Elfhag, K. and Rossner, S. (2005) Who succeeds in maintaining weight loss? A conceptual review of factors associated with weight loss maintenance and weight regain. *Obesity Reviews*. 6 (1), 67-85.
11. Foster, G.D., Makris, A.P. and Bailer, B.A. (2005) Behavioral treatment of obesity. *American Journal of Clinical Nutrition*. 82 (1), 230-235.

12. Goldstein LB. (2010). Physical activity and the risk of stroke. *Expert Rev Neurother*, 10(8), 1263-1265.
13. Hagberg, L.A., Lindahl, B., Nyberg, L. and Hellenius, M.L. (2008) Importance of enjoyment when promoting physical activity. *Scandinavian Journal of Medicine and Science in Sports*. 19 (5), 740-747.
14. Haus, G., Hoerr, S.L., Mavis, B. and Robison, J. (1994) Key modifiable factors in weight maintenance: Fat intake, exercise and weight cycling. *Journal of the Academy of Nutrition and Dietetics*. 94 (4), 409-413.
15. Jensen, P., Zachariae, C., Christensen, R., Geiker, N., Schaadt, B., Stender, S., Hansen, P., Astrup, A. & Skov, L. (2013) Effect of Weight Loss on the Severity of Psoriasis. *JAMA Dermatology*. 149 (7), 795.
16. Jolly, K., Lewis, A., Beach, J., Denley, J., Adab, P., Deeks, J., Daley, A. and Aveyard, P. (2011). Comparison of range of commercial or primary care led weight reduction programmes with minimal intervention control for weight loss in obesity: Lighten Up randomised controlled trial. *BMJ*. 343 (2), 6500.
17. Kahkoska, A., Watts, M., Driscoll, K., Bishop, F., Mihos, P., Thomas, J., Law, J., Jain, N. & Mayer-Davis, E. (2018). Understanding antagonism and synergism: A qualitative assessment of weight management in youth with Type 1 diabetes mellitus. *Obesity Medicine*, 9, 21-31.
18. Karfopoulou, E., Anastasiou, C.A., Avgeraki, E., Kosmidis, M.H. and Yannakoulia, M. (2016) The role of social support in weight loss maintenance: results from the MedWeight study. *Journal of Behavioural Medicine*. 39 (3), 511-518.
19. Klem, M.L., Wing, R.R., McGuire, M.T., Seagle, H.M. and Hill, J.O. (1997) A descriptive study of individuals successful at long-term maintenance of substantial weight loss. *The American Journal of Clinical Nutrition*. 66 (2), 39-46.
20. Klohe-Lehman, D., Freeland-Graves, J., Anderson, E., McDowell, T., Clarke, K., Hanss-Nuss, H., Cai, G., Puri, D. and Milani, T. (2006). Nutrition Knowledge Is Associated with Greater Weight

- Loss in Obese and Overweight Low-Income Mothers. *Journal of the American Dietetic Association*. 106 (1), 65-75.
21. Kruger, J., Blanck, H.M. and Gillespie, C. (2006) Dietary and physical activity behaviors among adults successful at weight loss maintenance. *International Journal of Behavioral Nutrition and Physical Activity*. 3 (17).
  22. Kruger, J., Galuska, D.A., Serdula, M.K. and Jones, D.A. (2004) Attempting to lose weight. Specific practices among U.S. adults. *The American College of Preventive Medicine*. 26 (5), 402-406.
  23. LaMonte MJ, Blair SN, Church TS. (2005). Physical activity and diabetes prevention. *Journal of applied physiology*, 99(3):1205-1213.
  24. Leahey, T.M., Xu, X., Unick, J.L. and Wing, R.R. (2014) A preliminary investigation of the role of self-control in behavioral weight loss treatment. *Obesity Research and Clinical Practice*. 8 (2), 49-53.
  25. Lemstra, M.L., Bird, Y., Nwankwo, C., Rogers, M. and Moraros, J. (2016) Weight loss intervention adherence and factors promoting adherence: a meta-analysis. *Patient Preference and Adherence*. 10 (1), 1547-1559.
  26. Li J, Siegrist J. (2012). Physical activity and risk of cardiovascular disease--a meta-analysis of prospective cohort studies. *International journal of environmental research and public health*, 9(2), 391-407.
  27. Linde, J., Jeffery, R., French, S., Pronk, N. and Boyle, R. (2005). Self-weighing in weight gain prevention and weight loss trials. *Annals of Behavioral Medicine*. 30 (3), 210-216.
  28. Mackintosh KA, Knowles ZR, Ridgers ND, Fairclough SJ. (2011). Using formative research to develop CHANGE!: a curriculum-based physical activity promoting intervention. *BMC Public Health*, 11,831, doi:10.1186/1471-2458-11-831.
  29. Martinsen EW. (2008). Physical activity in the prevention and treatment of anxiety and depression. *Nordic journal of psychiatry*, 62 Suppl 47:25-29

30. Metzgar, C.J., Preston, A.G., Miller, D.L. and Nickols-Richardson, S.M. (2014) Facilitators and barriers to weight loss and weight loss maintenance: A qualitative exploration. *Journal of Human Nutrition and Dietetics*. 28 (6), 593–603.
31. Montesi, L., El Ghoch, M., Brodosi, L., Calugi, S., Marchesini, G. and Dalle Grave, R. (2016) Long-term weight loss maintenance for obesity: a multidisciplinary approach. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. 9, 37-46.
32. Peters PG, Alessio HM, Hagerman AE, Ashton T, Nagy S, Wiley RL. (2006). Short-term isometric exercise reduces systolic blood pressure in hypertensive adults: possible role of reactive oxygen species. *International journal of cardiology*, 110(2):199-205.
33. Ridgers ND, Knowles ZR, Sayers J. (2012). Encouraging play in the natural environment: a child-focused case study of Forest School. *Child Geogr*, 10(1), 49-65. [doi:10.1080/14733285.2011.638176]
34. Ross, R., Dagnone, D., Jones, P.J., Smith, H., Paddags, A., Hudson, R. and Janssen, I. (2000) Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Annals of Internal Medicine*. 133 (2), 92-103.
35. Ryan, R.M., Frederick, C.M., Lepes, D., Rubio, N. and Sheldon, K.M. (1997) Intrinsic motivation and exercise adherence. *International Journal of Sport Psychology*. 28 (4), 335-354.
36. Saunders TJ, Gray CE, Poitras VJ, Chaput JP, Janssen I, Katzmarzyk PT, Olds T, Connor Gorber S, Kho ME, Sampson M, Tremblay MS, Carson V. (2016). Combinations of physical activity, sedentary behaviour and sleep: relationships with health indicators in school-aged children and youth. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*, 41(6 Suppl 3), S283-293.
37. Schifter, D. and Ajzen, I. (1985). Intention, perceived control, and weight loss: An application of the theory of planned behavior. *Journal of Personality and Social Psychology*. 49 (3), 843-851.

38. Shiri R, Solovieva S, Husgafvel-Pursiainen K, Telama R, Yang X, Viikari J, Raitakari OT, Viikari-Juntura E. (2013). The role of obesity and physical activity in non-specific and radiating low back pain: the Young Finns study. *Seminars in arthritis and rheumatism*, 42(6):640-650.
39. Spirduto, W.A., Thompson, H.S. and O'Brien, R.M. (1986) The effect of target behavior monitoring on weight loss and completion rate in behaviour modification program for weight reduction. *Addictive Behaviour*. 11 (3), 337-340.
40. Teixeira, P.J., Goings, S.B., Sardinha, L.B. and Lohman, T.G. (2005). A review of psychosocial pre-treatment predictors of weight control. *Obesity Reviews*. 6 (1), 43–65.
41. Telama R, Yang X, Leskinen E, Kankaanpaa A, Hirvensalo M, Tammelin T, Viikari JS, Raitakari OT. (2013). Tracking of Physical Activity from Early Childhood through Youth into Adulthood. *Med Sci Spor Exerc*, 46 (5), 955–962
42. Trost, S.G., Owen, N., Bauman, A.E., Sallis, J.F. and Brown, W. (2002) Correlates of adults' participation in physical activity: review and update. *Medicine and Science in Sports and Exercise*. 34 (8), pp. 1996–2001.
43. Wadden, T., Butryn, M. and Wilson, C. (2007). Lifestyle Modification for the Management of Obesity. *Gastroenterology*. 132(6), 2226-2238.
44. Wadden, T., Neiberg, R., Wing, R., Clark, J., Delahanty, L., Hill, J., Krakoff, J., Otto, A., Ryan, D. and Vitolins, M. (2011). Four-Year Weight Losses in the Look AHEAD Study: Factors Associated with Long-Term Success. *Obesity*. 19 (10), pp.1987-1998.
45. Weight management. (2003). Washington, D.C.: National Academies Press.
46. Welch, N., McNaughton, S.A., Hunter, W., Hume, C. and Crawford, D. (2009) Is the perception of time pressure a barrier to healthy eating and physical activity among women? *Public Health Nutrition*. 12 (7), 888-895.
47. WHO. (2010). Global recommendations on physical activity for health.
48. Wing, R.R. and Hill, J.O. (2001) Successful weight loss maintenance. *Annual Review of Nutrition*. 21 (1), 323–341.

49. Wolin KY, Yan Y, Colditz GA, Lee IM. (2009). Physical activity and colon cancer prevention: a meta-analysis. *British journal of cancer*, 100(4):611-616.
50. Wu, T., Gao, X., Chen, M. and van Dam, R. (2009). Long-term effectiveness of diet-plus-exercise interventions vs. diet-only interventions for weight loss: a meta-analysis. *Obesity Reviews*. 10 (3), 313-323.

1 **Figure captions**

2

3 FIGURE 1. Barriers to weight loss success for group and individual strategies

4

5 FIGURE 2. Contributors to weight loss for group strategies

6

7 FIGURE 3. Contributors to weight loss success for individual strategies

8

9 FIGURE 4. Strategy choice influences for group and individual weight loss

10


11

12

13


14

15




**FIGURE 1. Barriers to weight loss success for group and individual strategies**


**FIGURE 2.**  
**Contributors to weight loss for group strategies**





**FIGURE 3.**  
Contributors to weight loss success for individual strategies





**FIGURE 4. Strategy choice influences for group and individual weight loss**

